

| | AVE | TOTAL MAX | SURGEON STD | MODE | TTL | AVE | TEACHING ASSISTANT MAX | STD | MODE | TTL |
|---|------|--------------|----------------|------|------|-----|---------------------------|-----|------|-----|
| Skin/Soft Tissue Musculoskeletal | 31.1 | 78 | 22 | 12 | 839 | 8.3 | 139 | 27 | 0 | 224 |
| Burn debridement or grafting | 8.7 | 50 | 15 | 0 | 235 | 5.1 | 103 | 20 | 0 | 137 |
| Complex wound closure | 4.1 | 12 | 4 | 3 | 112 | 0.4 | 4 | 1 | 0 | 11 |
| Subcutaneous mastectomy | 1.0 | 5 | 1 | 0 | 28 | 0.1 | 1 | 0 | 0 | 4 |
| Pilonidal cyst excision | 3.3 | 12 | 3 | 1 | 88 | 0.3 | 3 | 1 | 0 | 9 |
| Major excision soft tissue tumor | 8.7 | 23 | 7 | 10 | 234 | 0.9 | 14 | 3 | 0 | 25 |
| Major soft tissue repair for trauma | 2.7 | 12 | 3 | 1 | 73 | 0.9 | 7 | 2 | 0 | 25 |
| Other | 2.6 | 19 | 4 | 0 | 69 | 0.5 | 10 | 2 | 0 | 13 |
| Head & Neck | 27.1 | 61 | 13 | 15 | 732 | 4.3 | 68 | 13 | 0 | 117 |
| Cleft lip/palate repair | 0.0 | 1 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 |
| Thyroidectomy (any) | 4.1 | 11 | 3 | 1 | 112 | 0.4 | 6 | 1 | 0 | 11 |
| Parathyroidectomy (any) | 0.2 | 2 | 1 | 0 | 5 | 0.0 | 1 | 0 | 0 | 1 |
| Branchial cleft cyst/sinus | 4.9 | 13 | 4 | 1 | 133 | 0.4 | 6 | 1 | 0 | 11 |
| Thyroglossal duct cyst/sinus | 4.0 | 9 | 2 | 4 | 107 | 0.2 | 3 | 1 | 0 | 5 |
| Cystic hygroma/lymphangioma | 2.9 | 8 | 2 | 4 | 77 | 0.2 | 2 | 1 | 0 | 6 |
| Dermoid/other cyst | 8.9 | 33 | 9 | 3 | 241 | 3.0 | 50 | 10 | 0 | 80 |
| Major tumor (head & neck) | 1.9 | 7 | 2 | 0 | 50 | 0.1 | 1 | 0 | 0 | 3 |
| Other | 0.2 | 3 | 1 | 0 | 6 | 0.0 | 0 | 0 | 0 | 0 |
| Thoracic | 67.1 | 116 | 24 | 62 | 1811 | 5.9 | 31 | 9 | 0 | 159 |
| Tracheostomy | 3.0 | 16 | 4 | 0 | 80 | 0.1 | 2 | 0 | 0 | 3 |
| Laryngeal or tracheal resection and/or reconstruction | 0.1 | 1 | 0 | 0 | 2 | 0.0 | 1 | 0 | 0 | 1 |
| Repair chest wall deformity | 12.7 | 61 | 13 | 4 | 342 | 1.0 | 7 | 2 | 0 | 26 |
| Resection chest wall tumor | 1.9 | 6 | 2 | 0 | 50 | 0.1 | 1 | 0 | 0 | 4 |
| Excision mediastinal cyst | 3.1 | 10 | 2 | 1 | 84 | 0.2 | 2 | 0 | 0 | 5 |
| Excision mediastinal tumor | 3.9 | 13 | 3 | 2 | 105 | 0.2 | 3 | 1 | 0 | 5 |
| Pulmonary resection tumor, congenital malformation, infection, et | 11.2 | 36 | 6 | 8 | 303 | 0.7 | 9 | 2 | 0 | 20 |
| Thoracotomy for trauma | 0.8 | 4 | 1 | 0 | 22 | 0.0 | 1 | 0 | 0 | 1 |
| Lung biopsy: open | 2.1 | 9 | 2 | 0 | 57 | 0.1 | 1 | 0 | 0 | 4 |
| Lung biopsy: scope | 5.9 | 29 | 6 | 5 | 158 | 0.6 | 5 | 1 | 0 | 16 |
| Decortication/pleurectomy/blebectomy: open | 1.1 | 6 | 1 | 0 | 30 | 0.2 | 1 | 0 | 0 | 5 |
| Decortication/pleurectomy/blebectomy: scope | 9.9 | 34 | 7 | 6 | 267 | 1.6 | 12 | 3 | 0 | 42 |
| Esophageal resection or replacement | 1.8 | 5 | 1 | 1 | 49 | 0.2 | 2 | 1 | 0 | 6 |
| Esophagomyotomy | 0.9 | 5 | 1 | 0 | 25 | 0.1 | 1 | 0 | 0 | 2 |
| Repair esophageal atresia and/or tracheoesophageal fistula | 8.6 | 15 | 4 | 4 | 231 | 0.7 | 8 | 2 | 0 | 18 |
| Other | 0.2 | 2 | 1 | 0 | 6 | 0.0 | 1 | 0 | 0 | 1 |
| Diaphragm | 18.4 | 37 | 8 | 19 | 498 | 1.8 | 25 | 5 | 0 | 49 |
| Repair diaphragmatic hernia | 12.9 | 21 | 4 | 15 | 347 | 0.8 | 5 | 1 | 0 | 21 |
| Plication of diaphragm | 1.0 | 3 | 1 | 1 | 28 | 0.2 | 5 | 1 | 0 | 5 |
| Transthoracic and/or retroperitoneal exposure for scoliosis | 4.4 | 22 | 5 | 1 | 118 | 0.9 | 17 | 3 | 0 | 23 |
| Other | 0.2 | 1 | 0 | 0 | 5 | 0.0 | 0 | 0 | 0 | 0 |

* AVE: Average number of operations performed by all residents MAX: Maximun number of operations performed by any resident
 STD: Standard deviation MDE: Mode, most common number of operations performed TTL: Total number of operations performed by all residents

| | AVE | TOTAL MAX | SURGEON STD | MODE | TTL | AVE | TEACHING ASSISTANT MAX | STD | MODE | TTL |
|---|-------|--------------|----------------|------|------|------|---------------------------|-----|------|------|
| Cardiovascular | 113.2 | 300 | 57 | . | 3056 | 21.3 | 214 | 44 | 0 | 574 |
| Patent ductus arteriosus | 4.0 | 26 | 7 | 0 | 107 | 0.0 | 0 | 0 | 0 | 0 |
| Coarctation | 0.1 | 2 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 |
| Vascular ring | 0.1 | 2 | 0 | 0 | 4 | 0.0 | 1 | 0 | 0 | 1 |
| Any open heart procedure | 0.0 | 1 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 |
| Any closed heart procedure | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| Aortopexy | 0.5 | 3 | 1 | 0 | 13 | 0.0 | 1 | 0 | 0 | 1 |
| Renal artery reconstruction | 0.1 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 |
| Peripheral artery reconstruction | 0.6 | 4 | 1 | 0 | 15 | 0.0 | 1 | 0 | 0 | 1 |
| Construction or take down AV fistula/shunt | 1.1 | 7 | 2 | 0 | 30 | 0.1 | 1 | 0 | 0 | 3 |
| Surgical placement/removal central access line | 89.9 | 262 | 56 | . | 2426 | 19.5 | 210 | 43 | 0 | 526 |
| Dialysis access insertion/removal | 2.1 | 14 | 3 | 0 | 56 | 0.4 | 6 | 1 | 0 | 12 |
| Cannulate/Decannulate ECMO | 14.1 | 29 | 8 | 12 | 380 | 1.1 | 8 | 2 | 0 | 29 |
| Major vessel reconstruction | 0.7 | 4 | 1 | 0 | 20 | 0.0 | 1 | 0 | 0 | 1 |
| Other | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| Abdominal | 348.5 | 665 | 122 | 276 | 9410 | 38.3 | 226 | 49 | 0 | 1034 |
| Antireflux procedure: open | 22.1 | 84 | 21 | 10 | 596 | 1.7 | 20 | 4 | 0 | 46 |
| Antireflux procedure: scope | 25.8 | 83 | 23 | 12 | 696 | 1.1 | 8 | 2 | 0 | 30 |
| Pyloroplasty/gastric resection w/wout vagotomy | 2.0 | 13 | 3 | 1 | 54 | 0.0 | 1 | 0 | 0 | 1 |
| Any gastrostomy/jejunostomy: open | 11.2 | 64 | 12 | 9 | 303 | 0.8 | 5 | 1 | 0 | 22 |
| Any gastrostomy/jejunostomy: scope | 7.0 | 41 | 9 | 0 | 190 | 1.2 | 12 | 3 | 0 | 33 |
| Pyloromyotomy | 32.5 | 72 | 16 | 19 | 877 | 4.5 | 22 | 5 | 1 | 122 |
| Operation for malrotation | 12.9 | 26 | 6 | 10 | 348 | 0.9 | 5 | 1 | 0 | 25 |
| Duodenal atresia | 4.1 | 9 | 3 | 7 | 110 | 0.3 | 2 | 1 | 0 | 7 |
| Repair intestinal atresia, stenosis or web | 10.8 | 26 | 6 | 11 | 292 | 0.4 | 2 | 1 | 0 | 10 |
| Intestinal resection/repair or ostomy for necrotizing enterocolit | 21.3 | 38 | 8 | 12 | 576 | 1.4 | 8 | 2 | 0 | 37 |
| Intestinal resection/repair or ostomy for inflammatory bowel dise | 4.1 | 11 | 3 | 5 | 111 | 0.3 | 4 | 1 | 0 | 9 |
| Intestinal resection/repair or ostomy for trauma | 4.5 | 17 | 4 | 3 | 121 | 0.6 | 3 | 1 | 0 | 16 |
| Intestinal resection (Meckel's, duplication, meconium ileus, etc. | 8.5 | 19 | 4 | 3 | 229 | 0.7 | 7 | 2 | 0 | 18 |
| Bowel resection for Crohn's disease | 2.5 | 8 | 2 | 1 | 67 | 0.2 | 2 | 1 | 0 | 6 |
| Laparotomy or resection for intussusception | 4.4 | 10 | 2 | 5 | 119 | 0.5 | 4 | 1 | 0 | 14 |
| Ostomy for anorectal malformation | 4.0 | 11 | 3 | 3 | 109 | 0.2 | 2 | 1 | 0 | 6 |
| Ostomy for Hirschsprung's | 5.5 | 15 | 4 | 5 | 149 | 0.3 | 2 | 1 | 0 | 7 |
| Ostomy for Other | 1.8 | 8 | 2 | 0 | 49 | 0.1 | 1 | 0 | 0 | 4 |
| Closure/revision any ostomy | 18.3 | 39 | 10 | 31 | 494 | 1.7 | 15 | 3 | 0 | 47 |
| Appendectomy: open | 14.0 | 80 | 16 | 8 | 379 | 4.4 | 26 | 8 | 0 | 119 |
| Appendectomy: scope | 38.1 | 108 | 25 | 41 | 1028 | 10.4 | 65 | 17 | 1 | 281 |
| Perineal procedure for imperforate anus | 13.4 | 42 | 8 | 13 | 361 | 0.6 | 6 | 1 | 0 | 15 |
| Pull through for Hirschsprung's: open | 7.3 | 17 | 4 | 5 | 197 | 0.2 | 4 | 1 | 0 | 6 |
| Pull through for Hirschsprung's: scope | 2.8 | 8 | 2 | 0 | 76 | 0.0 | 1 | 0 | 0 | 1 |
| Pull through for IBD or polyposis: open | 2.3 | 7 | 2 | 2 | 61 | 0.2 | 3 | 1 | 0 | 6 |
| Pull through for IBD or polyposis: scope | 0.9 | 5 | 1 | 0 | 24 | 0.1 | 1 | 0 | 0 | 2 |
| Exploratory laparotomy with or without biopsy | 16.7 | 37 | 9 | 12 | 451 | 1.5 | 12 | 3 | 0 | 40 |
| Excision of omental/mesenteric cyst | 1.7 | 6 | 2 | 0 | 47 | 0.2 | 4 | 1 | 0 | 6 |
| Omphalocele (any surgical repair) | 5.0 | 11 | 3 | 4 | 134 | 0.3 | 2 | 1 | 0 | 9 |
| Gastroschisis (any surgical repair) | 14.2 | 56 | 12 | 3 | 384 | 1.4 | 7 | 2 | 0 | 37 |
| Resection urachal remnant | 2.5 | 6 | 2 | 2 | 68 | 0.1 | 2 | 0 | 0 | 3 |
| Resection omphalomesenteric duct/cyst | 0.9 | 4 | 1 | 0 | 23 | 0.1 | 1 | 0 | 0 | 2 |

* AVE: Average number of operations performed by all residents MAX: Maximum number of operations performed by any resident
 STD: Standard deviation MDE: Mode, most common number of operations performed TTL: Total number of operations performed by all residents

| | AVE | TOTAL MAX | SURGEON STD | MODE | TTL | AVE | TEACHING ASSISTANT MAX | STD | MODE | TTL |
|--|--------------|--------------|----------------|-----------|-------------|-------------|---------------------------|-----------|----------|------------|
| Excision neuroblastoma/adrenal/other retroperitoneal tumor | 10.9 | 26 | 6 | 6 | 294 | 0.3 | 3 | 1 | 0 | 9 |
| Excision sacrococcygeal teratoma | 1.8 | 5 | 1 | 2 | 49 | 0.3 | 4 | 1 | 0 | 8 |
| Diagnostic laparoscopy | 5.5 | 26 | 5 | 5 | 149 | 0.5 | 3 | 1 | 0 | 14 |
| Other | 7.2 | 25 | 7 | 3 | 195 | 0.6 | 9 | 2 | 0 | 16 |
| Hernia Repair | 109.2 | 215 | 50 | 80 | 2948 | 20.5 | 253 | 52 | 0 | 553 |
| Ped repair inguinal hernia (uni/bilateral) | 43.2 | 101 | 27 | 21 | 1166 | 10.6 | 148 | 30 | 1 | 285 |
| Infant (<6 mos of age) rep inguinal hernia (uni/bilateral) | 51.0 | 103 | 26 | 47 | 1378 | 3.1 | 32 | 7 | 0 | 84 |
| Repair umbilical hernia | 9.6 | 34 | 8 | 2 | 260 | 5.5 | 59 | 14 | 0 | 148 |
| Repair ventral hernia | 5.1 | 15 | 4 | 3 | 139 | 1.3 | 15 | 4 | 0 | 36 |
| Other | 0.2 | 1 | 0 | 0 | 5 | 0.0 | 0 | 0 | 0 | 0 |
| Liver/Biliary | 44.6 | 75 | 16 | 29 | 1204 | 5.5 | 29 | 8 | 0 | 149 |
| Major hepatic resection/repair: tumor | 2.5 | 6 | 2 | 3 | 67 | 0.2 | 1 | 0 | 0 | 5 |
| Major hepatic resection/repair: trauma | 0.6 | 4 | 1 | 0 | 15 | 0.0 | 1 | 0 | 0 | 1 |
| Major hepatic resection/repair: other | 0.1 | 1 | 0 | 0 | 4 | 0.0 | 1 | 0 | 0 | 1 |
| Lysis of adhesions | 4.3 | 13 | 4 | 0 | 115 | 0.4 | 4 | 1 | 0 | 12 |
| Liver biopsy: open | 3.9 | 11 | 3 | 3 | 105 | 0.4 | 5 | 1 | 0 | 10 |
| Liver biopsy: scope | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| Liver harvest | 0.2 | 5 | 1 | 0 | 6 | 0.0 | 0 | 0 | 0 | 0 |
| Liver transplant | 1.1 | 12 | 3 | 0 | 30 | 0.1 | 2 | 0 | 0 | 2 |
| Cholecystectomy with/without common bile duct exploration: open | 1.4 | 7 | 2 | 0 | 39 | 0.2 | 1 | 0 | 0 | 6 |
| Cholecystectomy with/without common bile duct exploration: scope | 12.1 | 31 | 7 | 7 | 326 | 3.0 | 15 | 4 | 0 | 80 |
| Portoenterostomy | 2.9 | 7 | 2 | 4 | 77 | 0.1 | 2 | 0 | 0 | 4 |
| Excision choledochal cyst | 2.5 | 6 | 1 | 1 | 67 | 0.1 | 1 | 0 | 0 | 2 |
| Portosystemic shunts or other operations for portal hypertension | 1.1 | 18 | 4 | 0 | 31 | 0.1 | 1 | 0 | 0 | 2 |
| Pancreatic resection for: trauma | 0.4 | 3 | 1 | 0 | 12 | 0.0 | 1 | 0 | 0 | 1 |
| Pancreatic resection for: hyperinsulinism | 1.1 | 21 | 4 | 0 | 29 | 0.0 | 0 | 0 | 0 | 0 |
| Pancreatic resection for: tumor | 0.8 | 4 | 1 | 0 | 22 | 0.1 | 1 | 0 | 0 | 3 |
| Operations for pseudocyst | 0.7 | 3 | 1 | 0 | 18 | 0.0 | 1 | 0 | 0 | 1 |
| Splenorrhaphy | 0.1 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 |
| Splenectomy: open | 2.0 | 6 | 2 | 1 | 55 | 0.1 | 2 | 0 | 0 | 4 |
| Splenectomy: scope | 6.8 | 19 | 4 | 3 | 184 | 0.6 | 5 | 1 | 0 | 15 |
| Other | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| Genitourinary | 65.1 | 164 | 30 | 45 | 1758 | 7.2 | 82 | 17 | 0 | 194 |
| Nephrectomy (total or partial): tumor | 7.1 | 17 | 4 | 6 | 193 | 0.4 | 5 | 1 | 0 | 12 |
| Nephrectomy (total or partial): trauma | 0.1 | 2 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 0 |
| Nephrectomy (total or partial): cystic dysplasia | 0.2 | 3 | 1 | 0 | 6 | 0.1 | 1 | 0 | 0 | 2 |
| Nephrectomy (total or partial): other | 0.1 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 0 |
| Renal transplant | 2.7 | 15 | 5 | 0 | 72 | 0.2 | 4 | 1 | 0 | 6 |
| Renal biopsy (open) | 0.9 | 3 | 1 | 0 | 24 | 0.1 | 1 | 0 | 0 | 2 |
| Cystectomy (partial or total) | 0.2 | 2 | 1 | 0 | 6 | 0.0 | 0 | 0 | 0 | 0 |
| Nephrostomy | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| Operation for nephro-uretero lithiasis | 0.4 | 3 | 1 | 0 | 10 | 0.0 | 0 | 0 | 0 | 0 |
| Enteric conduit (any) | 0.3 | 4 | 1 | 0 | 8 | 0.0 | 0 | 0 | 0 | 0 |
| Bladder augmentation or repair (any) | 0.5 | 4 | 1 | 0 | 14 | 0.0 | 1 | 0 | 0 | 1 |
| Urinary undiversion | 0.0 | 1 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 |
| Pyeloplasty/UPJ reconstruction | 0.9 | 6 | 1 | 0 | 23 | 0.0 | 1 | 0 | 0 | 1 |
| Ureteral reconstruction/reimplantation | 1.4 | 6 | 2 | 0 | 37 | 0.0 | 1 | 0 | 0 | 1 |
| Reconstruct cloacal extrophy | 1.8 | 15 | 3 | 0 | 49 | 0.0 | 0 | 0 | 0 | 0 |
| Circumcision (OR only) | 12.6 | 52 | 13 | 4 | 340 | 4.0 | 50 | 12 | 0 | 109 |

* AVE: Average number of operations performed by all residents MAX: Maximum number of operations performed by any resident
 STD: Standard deviation MDE: Mode, most common number of operations performed TTL: Total number of operations performed by all residents

| | SURGEON | | | | | TEACHING ASSISTANT | | | | |
|---|--------------|-------------|------------|-------------|--------------|--------------------|------------|------------|----------|-------------|
| | AVE | MAX | STD | MODE | TTL | AVE | MAX | STD | MODE | TTL |
| Orchidopexy: open | 16.4 | 42 | 8 | 10 | 443 | 1.0 | 15 | 3 | 0 | 28 |
| Orchidopexy: scope | 2.1 | 13 | 3 | 0 | 57 | 0.1 | 1 | 0 | 0 | 3 |
| Orchiectomy | 1.1 | 7 | 2 | 0 | 30 | 0.0 | 1 | 0 | 0 | 1 |
| Operation for torsion testis or appendages | 0.5 | 3 | 1 | 0 | 13 | 0.1 | 1 | 0 | 0 | 2 |
| Operation for varicocele: open | 0.1 | 1 | 0 | 0 | 4 | 0.1 | 2 | 0 | 0 | 2 |
| Operation for varicocele: scope | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| Procedures for intersex (vaginal reconstruction, clitoroplasty, e1.0) | 1.0 | 5 | 1 | 0 | 26 | 0.0 | 0 | 0 | 0 | 0 |
| Oophorectomy (partial or total) | 8.4 | 19 | 4 | 6 | 227 | 0.6 | 5 | 1 | 0 | 16 |
| Hysterectomy/salpingectomy | 0.7 | 3 | 1 | 0 | 18 | 0.1 | 1 | 0 | 0 | 2 |
| Repair complex laceration vagina/perineum | 1.3 | 5 | 1 | 0 | 35 | 0.1 | 1 | 0 | 0 | 3 |
| Other | 4.3 | 32 | 8 | 0 | 116 | 0.1 | 2 | 0 | 0 | 3 |
| Endoscopic Procedures | 52.9 | 130 | 28 | 42 | 1429 | 4.6 | 58 | 11 | 0 | 123 |
| Diagnostic thoracoscopy | 0.6 | 3 | 1 | 0 | 15 | 0.0 | 1 | 0 | 0 | 1 |
| Diagnostic laparoscopy | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| Cystoscopy | 3.1 | 35 | 7 | 0 | 84 | 0.0 | 1 | 0 | 0 | 1 |
| Bronchoscopy | 9.3 | 61 | 12 | 3 | 251 | 0.4 | 3 | 1 | 0 | 10 |
| Esophagoscopy | 6.0 | 18 | 4 | 3 | 162 | 1.0 | 18 | 4 | 0 | 26 |
| Removal foreign body esophagus or trachea | 16.7 | 57 | 16 | 1 | 451 | 1.4 | 10 | 3 | 0 | 39 |
| Esophageal dilatation | 13.0 | 35 | 8 | 8 | 350 | 1.4 | 23 | 5 | 0 | 37 |
| Colonoscopy | 0.9 | 7 | 2 | 0 | 23 | 0.0 | 1 | 0 | 0 | 1 |
| Sigmoidoscopy | 3.3 | 11 | 3 | 1 | 89 | 0.3 | 6 | 1 | 0 | 8 |
| Other endoscopy | 105.8 | 223 | 50 | 90 | 2857 | 5.2 | 113 | 22 | 0 | 141 |
| Trauma | 105.8 | 223 | 50 | 90 | 2857 | 5.2 | 113 | 22 | 0 | 141 |
| ===== | | | | | | | | | | |
| Total | 983.0 | 1644 | 255 | 1135 | 26542 | 122.9 | 968 | 213 | 6 | 3317 |
| Defined Categories | | | | | | | | | | |
| Tumors | 48.0 | 83 | 17 | 38 | 1297 | 3.1 | 31 | 6 | 0 | 85 |
| Important Pediatric Surgical Cases | 71.8 | 162 | 28 | 63 | 1938 | 4.1 | 42 | 9 | 0 | 110 |
| Neonate | 97.0 | 182 | 31 | 95 | 2619 | 6.6 | 38 | 10 | 0 | 178 |

* AVE: Average number of operations performed by all residents MAX: Maximum number of operations performed by any resident
 STD: Standard deviation MDE: Mode, most common number of operations performed TTL: Total number of operations performed by all residents