



Accreditation Council for Graduate Medical Education

Pediatric Surgery Case Logs

National Data Report

Prepared by: Department of Applications and Data Analysis

- I. National Resident Statistics Main Table
 - A. This main table provides a national picture of educational experiences. The report is ideal for establishing and assessing national policy related to resident practical experiences. Data is broken out by resident role.
 - B. Descriptive statistics have been carefully chosen to provide the most useful information for judging performance at that national level.
 - i. "Natl Res AVE" - The national average describes the mean number of procedures performed per resident in the nation.
 - ii. "Natl Res STD" - The standard deviation indicates how tightly the scores are clustered around the mean in the set of data.
 - iii. "Natl Res MIN" - The minimum value among procedure counts in the nation.
 - iv. "Natl Res MED" - The median value among procedure counts in the nation.
 - v. "Natl Res MAX" - The maximum value seen among procedure counts in the nation.

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = All Roles

[PAGE 1] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|----------------------------------|--------------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Skin/Soft Tissue/Musculoskeletal | Burn debridement or grafting | 9.9 | 23 | 0 | 1 | 121 | 3.9 | 17 | 0 | 0 | 106 | 13.8 | 39 | 0 | 1 | 227 |
| | Complex wound closure | 4.7 | 6 | 0 | 4 | 35 | 1.1 | 2 | 0 | 0 | 8 | 5.7 | 7 | 0 | 4 | 36 |
| | Breast | 3.4 | 3 | 0 | 3 | 15 | 0.1 | 0 | 0 | 0 | 1 | 3.5 | 3 | 0 | 3 | 15 |
| | Pilonidal cyst excision | 3.6 | 4 | 0 | 3 | 20 | 0.0 | 0 | 0 | 0 | 0 | 3.6 | 4 | 0 | 3 | 20 |
| | Major excision soft tissue tumor | 8.3 | 4 | 1 | 7 | 22 | 0.4 | 1 | 0 | 0 | 2 | 8.6 | 4 | 1 | 8 | 22 |
| | Major soft tissue repair for trauma | 4.8 | 6 | 0 | 2 | 25 | 0.6 | 2 | 0 | 0 | 8 | 5.5 | 8 | 0 | 2 | 30 |
| | Other (skin/soft tissue/musc) | 11.0 | 8 | 0 | 9 | 36 | 1.1 | 1 | 0 | 1 | 5 | 12.1 | 8 | 0 | 11 | 40 |
| | TOTAL - Skin/Soft Tissue/Musc | 45.6 | 35 | 10 | 37 | 190 | 7.2 | 20 | 0 | 3 | 124 | 52.9 | 53 | 10 | 41 | 314 |
| Head & Neck | Thyroidectomy (any) | 5.8 | 7 | 0 | 3 | 23 | 0.1 | 0 | 0 | 0 | 1 | 5.9 | 7 | 0 | 3 | 23 |
| | Parathyroidectomy (any) | 0.6 | 1 | 0 | 0 | 4 | 0.0 | 0 | 0 | 0 | 0 | 0.6 | 1 | 0 | 0 | 4 |
| | Branchial cleft cyst/sinus | 3.6 | 2 | 0 | 4 | 7 | 0.1 | 0 | 0 | 0 | 1 | 3.6 | 2 | 0 | 4 | 7 |
| | Thyroglossal duct cyst/sinus | 2.6 | 2 | 0 | 2 | 14 | 0.0 | 0 | 0 | 0 | 0 | 2.6 | 2 | 0 | 2 | 14 |
| | Cystic hygroma/lymphangioma | 0.7 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 1 | 0.8 | 1 | 0 | 0 | 3 |
| | Dermoid/other cyst | 7.7 | 7 | 0 | 5 | 31 | 0.6 | 2 | 0 | 0 | 8 | 8.3 | 7 | 0 | 6 | 31 |
| | Major tumor (head & neck) | 1.0 | 1 | 0 | 1 | 4 | 0.1 | 0 | 0 | 0 | 1 | 1.1 | 1 | 0 | 1 | 4 |
| | Other (head & neck) | 4.0 | 3 | 0 | 3 | 11 | 0.2 | 0 | 0 | 0 | 1 | 4.2 | 3 | 0 | 3 | 11 |
| | TOTAL - Head & Neck | 26.1 | 11 | 11 | 24 | 55 | 0.9 | 2 | 0 | 0 | 8 | 27.0 | 11 | 11 | 25 | 57 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = All Roles

[PAGE 2]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|----------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Thoracic | Tracheostomy | 1.9 | 3 | 0 | 0 | 11 | 0.3 | 1 | 0 | 0 | 2 | 2.2 | 3 | 0 | 1 | 12 |
| | Laryngeal or tracheal resect and/or recon | 0.2 | 1 | 0 | 0 | 2 | 0.1 | 1 | 0 | 0 | 4 | 0.3 | 1 | 0 | 0 | 6 |
| | Repair chest wall deformity | 13.9 | 9 | 2 | 13 | 50 | 0.2 | 0 | 0 | 0 | 2 | 14.1 | 9 | 2 | 13 | 52 |
| | Resection chest wall tumor | 1.5 | 1 | 0 | 1 | 5 | 0.1 | 0 | 0 | 0 | 2 | 1.6 | 1 | 0 | 1 | 5 |
| | Excision mediastinal cyst | 2.9 | 2 | 0 | 2 | 9 | 0.1 | 0 | 0 | 0 | 2 | 3.1 | 2 | 0 | 3 | 10 |
| | Excision mediastinal tumor | 3.2 | 2 | 0 | 3 | 10 | 0.4 | 1 | 0 | 0 | 2 | 3.5 | 3 | 0 | 4 | 10 |
| | Pulmonary resection: open | 8.2 | 7 | 1 | 7 | 40 | 0.7 | 1 | 0 | 0 | 8 | 8.8 | 7 | 1 | 7 | 40 |
| | Pulmonary resection: scope | 7.4 | 4 | 1 | 6 | 18 | 0.2 | 0 | 0 | 0 | 1 | 7.6 | 4 | 1 | 6 | 19 |
| | Thoracotomy for trauma | 0.7 | 1 | 0 | 1 | 3 | 0.1 | 0 | 0 | 0 | 2 | 0.8 | 1 | 0 | 1 | 3 |
| | Lung biopsy: open | 2.1 | 2 | 0 | 2 | 7 | 0.8 | 2 | 0 | 0 | 10 | 2.9 | 3 | 0 | 2 | 14 |
| | Lung biopsy: scope | 4.0 | 4 | 0 | 2 | 14 | 0.4 | 1 | 0 | 0 | 4 | 4.5 | 4 | 0 | 3 | 17 |
| | Decortication/pleurectomy/blebectomy: open | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.2 | 0 | 0 | 0 | 2 |
| | Decortication/pleurectomy/blebectomy: scope | 5.2 | 3 | 0 | 4 | 16 | 0.5 | 1 | 0 | 0 | 4 | 5.7 | 4 | 0 | 5 | 16 |
| | Esophageal resection or replacement | 2.6 | 2 | 0 | 2 | 10 | 0.2 | 0 | 0 | 0 | 2 | 2.8 | 2 | 0 | 2 | 11 |
| | Esophagomyotomy | 1.3 | 1 | 0 | 1 | 4 | 0.1 | 0 | 0 | 0 | 1 | 1.4 | 1 | 0 | 1 | 4 |
| | Repair esophageal atresia/tracheoesophageal fistula | 10.2 | 6 | 2 | 9 | 33 | 0.1 | 0 | 0 | 0 | 1 | 10.3 | 6 | 2 | 9 | 34 |
| | Thoracoscopy: other | 0.6 | 1 | 0 | 0 | 4 | 0.5 | 2 | 0 | 0 | 9 | 1.2 | 2 | 0 | 0 | 11 |
| | Thoracotomy: other | 1.3 | 2 | 0 | 1 | 11 | 0.5 | 1 | 0 | 0 | 8 | 1.8 | 3 | 0 | 1 | 13 |
| | Other (thoracic) | 1.1 | 3 | 0 | 0 | 14 | 0.4 | 1 | 0 | 0 | 5 | 1.5 | 3 | 0 | 0 | 14 |
| | TOTAL - Thoracic | 68.5 | 21 | 24 | 65 | 118 | 5.8 | 7 | 0 | 3 | 31 | 74.3 | 24 | 27 | 72 | 132 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = All Roles

[PAGE 3]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|----------------|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Diaphragm | Repair diaphragmatic hernia | 13.0 | 6 | 4 | 12 | 31 | 0.2 | 1 | 0 | 0 | 3 | 13.2 | 6 | 4 | 12 | 33 |
| | Plication of diaphragm | 0.8 | 1 | 0 | 1 | 2 | 0.1 | 0 | 0 | 0 | 1 | 0.9 | 1 | 0 | 1 | 3 |
| | Transthoracic and/or retroperitoneal exp for scoliosis | 0.8 | 2 | 0 | 0 | 6 | 0.1 | 0 | 0 | 0 | 2 | 0.9 | 2 | 0 | 0 | 7 |
| | Other (diaphragm) | 0.2 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 4 | 0.5 | 1 | 0 | 0 | 4 |
| | TOTAL - Diaphragm | 14.8 | 7 | 5 | 14 | 33 | 0.7 | 1 | 0 | 0 | 4 | 15.4 | 7 | 5 | 15 | 36 |
| Cardiovascular | Patent ductus arteriosus | 2.0 | 5 | 0 | 0 | 30 | 0.0 | 0 | 0 | 0 | 1 | 2.0 | 5 | 0 | 0 | 30 |
| | Vascular ring | 0.3 | 1 | 0 | 0 | 5 | 0.1 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 6 |
| | Any open heart procedure | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Any closed heart procedure | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 1 |
| | Aortopexy | 0.7 | 1 | 0 | 0 | 7 | 0.0 | 0 | 0 | 0 | 0 | 0.7 | 1 | 0 | 0 | 7 |
| | Renal artery reconstruction | 0.2 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 1 |
| | Peripheral artery reconstruction | 0.2 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 2 |
| | Construction or take down AV fistula/shunt | 0.5 | 1 | 0 | 0 | 6 | 0.0 | 0 | 0 | 0 | 0 | 0.5 | 1 | 0 | 0 | 6 |
| | Surgical placement/removal central access line | 79.5 | 37 | 32 | 77 | 191 | 16.9 | 17 | 0 | 14 | 76 | 96.4 | 46 | 37 | 94 | 267 |
| | Dialysis access insertion/removal | 1.2 | 2 | 0 | 1 | 10 | 0.1 | 0 | 0 | 0 | 2 | 1.3 | 2 | 0 | 1 | 10 |
| | Cannulate/ECMO | 15.6 | 8 | 0 | 15 | 37 | 0.2 | 1 | 0 | 0 | 2 | 15.8 | 8 | 0 | 15 | 37 |
| | Major vessel reconstruction | 1.3 | 2 | 0 | 0 | 13 | 0.2 | 1 | 0 | 0 | 3 | 1.5 | 3 | 0 | 1 | 16 |
| | Vascular trauma | 0.6 | 2 | 0 | 0 | 9 | 0.0 | 0 | 0 | 0 | 0 | 0.6 | 2 | 0 | 0 | 9 |
| | Vascular: post-operative exploration | 0.2 | 1 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 2 |
| | TOTAL - Cardiovascular | 102.3 | 40 | 47 | 95 | 239 | 17.7 | 18 | 0 | 14 | 79 | 119.9 | 51 | 54 | 111 | 318 |

Based upon the number of procedures logged by residents in the indicated role.

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = All Roles

[PAGE 4] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|-----------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Abdominal | Antireflux procedure: open | 7.4 | 9 | 0 | 6 | 42 | 0.9 | 1 | 0 | 1 | 4 | 8.3 | 9 | 0 | 6 | 43 |
| | Antireflux procedure: scope | 26.5 | 21 | 4 | 18 | 97 | 0.9 | 1 | 0 | 0 | 5 | 27.4 | 22 | 4 | 19 | 97 |
| | Pyloroplasty/gastric resection w/wout vagotomy | 2.4 | 2 | 0 | 2 | 10 | 0.6 | 1 | 0 | 0 | 3 | 3.0 | 3 | 0 | 2 | 10 |
| | Pyloroplasty/gastric resect w/wout vagotomy - lap | 1.7 | 5 | 0 | 0 | 29 | 0.1 | 0 | 0 | 0 | 2 | 1.9 | 5 | 0 | 0 | 29 |
| | Any gastrostomy/jejunostomy: open | 13.4 | 10 | 3 | 11 | 45 | 9.9 | 9 | 0 | 8 | 35 | 23.4 | 15 | 4 | 18 | 68 |
| | Any gastrostomy/jejunostomy: scope | 34.8 | 24 | 7 | 25 | 102 | 14.9 | 15 | 0 | 10 | 58 | 49.7 | 30 | 11 | 44 | 134 |
| | Pyloromyotomy: open | 8.3 | 11 | 0 | 3 | 45 | 0.1 | 0 | 0 | 0 | 1 | 8.3 | 11 | 0 | 3 | 45 |
| | Pyloromyotomy: scope | 31.4 | 21 | 0 | 27 | 109 | 0.2 | 0 | 0 | 0 | 1 | 31.5 | 21 | 0 | 27 | 109 |
| | Operation for malrotation | 10.7 | 4 | 1 | 11 | 20 | 1.4 | 2 | 0 | 1 | 9 | 12.1 | 5 | 1 | 13 | 24 |
| | Duodenal atresia | 5.4 | 3 | 0 | 5 | 13 | 0.1 | 0 | 0 | 0 | 2 | 5.5 | 3 | 0 | 6 | 13 |
| | Repair intestinal atresia, stenosis or web | 11.8 | 8 | 1 | 10 | 42 | 1.9 | 2 | 0 | 2 | 6 | 13.7 | 8 | 1 | 12 | 43 |
| | Intestinal resection: open | 23.7 | 11 | 6 | 21 | 52 | 5.9 | 6 | 0 | 4 | 26 | 29.6 | 13 | 9 | 30 | 53 |
| | Intestinal resection: scope | 2.6 | 2 | 0 | 2 | 8 | 0.1 | 0 | 0 | 0 | 1 | 2.6 | 2 | 0 | 2 | 8 |
| | Intestinal resection/repair or ostomy for IBD | 2.5 | 2 | 0 | 2 | 9 | 0.3 | 1 | 0 | 0 | 3 | 2.8 | 2 | 0 | 3 | 11 |
| | Intestinal resect/repair or ostomy for IBD - lap | 5.3 | 4 | 0 | 5 | 16 | 0.1 | 0 | 0 | 0 | 1 | 5.4 | 4 | 0 | 5 | 16 |
| | Intestinal resection/repair or ostomy for trauma | 4.6 | 4 | 0 | 4 | 16 | 1.9 | 2 | 0 | 1 | 8 | 6.5 | 5 | 0 | 5 | 18 |
| | Intestinal resect/repair or ostomy for trauma - lap | 0.4 | 1 | 0 | 0 | 4 | 0.0 | 0 | 0 | 0 | 1 | 0.5 | 1 | 0 | 0 | 4 |
| | Bowel resection: open | 4.5 | 5 | 0 | 4 | 29 | 0.7 | 1 | 0 | 0 | 5 | 5.3 | 5 | 0 | 4 | 29 |
| | Bowel resection: scope | 1.7 | 2 | 0 | 1 | 9 | 0.1 | 0 | 0 | 0 | 1 | 1.8 | 2 | 0 | 1 | 9 |
| | Lap/resec for intussusception-malrotation-volvulus | 7.3 | 3 | 2 | 7 | 14 | 0.4 | 1 | 0 | 0 | 4 | 7.7 | 3 | 2 | 7 | 15 |
| | Ostomy for anorectal malformation: open | 3.8 | 2 | 0 | 4 | 10 | 0.5 | 1 | 0 | 0 | 3 | 4.3 | 3 | 1 | 4 | 12 |
| | Ostomy for anorectal malformation: laparoscopic | 0.6 | 1 | 0 | 0 | 6 | 0.2 | 0 | 0 | 0 | 1 | 0.8 | 1 | 0 | 0 | 6 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = All Roles

[PAGE 5] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|-----------|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Abdominal | Ostomy for Hirschsprung's: open | 1.8 | 2 | 0 | 1 | 6 | 0.1 | 0 | 0 | 0 | 1 | 1.9 | 2 | 0 | 1 | 6 |
| | Ostomy for Hirschsprung's: laparoscopic | 1.3 | 1 | 0 | 1 | 8 | 0.1 | 0 | 0 | 0 | 2 | 1.4 | 2 | 0 | 1 | 8 |
| | Ostomy for other: open | 7.0 | 4 | 1 | 7 | 16 | 1.8 | 2 | 0 | 1 | 7 | 8.8 | 5 | 1 | 9 | 20 |
| | Ostomy for other: laparoscopic | 0.6 | 1 | 0 | 0 | 3 | 0.2 | 0 | 0 | 0 | 2 | 0.8 | 1 | 0 | 0 | 4 |
| | Closure/revision any ostomy/fistula: open | 25.0 | 10 | 6 | 25 | 55 | 2.9 | 3 | 0 | 2 | 11 | 27.9 | 11 | 6 | 28 | 62 |
| | Closure/revision any ostomy/fistula: scope | 0.4 | 1 | 0 | 0 | 7 | 0.0 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 8 |
| | Appendectomy: open | 3.3 | 5 | 0 | 2 | 25 | 3.8 | 6 | 0 | 1 | 26 | 7.1 | 8 | 0 | 3 | 34 |
| | Appendectomy: scope | 108.5 | 63 | 22 | 100 | 296 | 2.4 | 3 | 0 | 2 | 16 | 110.9 | 64 | 24 | 100 | 303 |
| | Perineal procedure for imperforate anus | 13.8 | 6 | 4 | 14 | 30 | 0.2 | 0 | 0 | 0 | 2 | 14.1 | 6 | 4 | 14 | 31 |
| | Pull through for Hirschsprung's: open | 5.1 | 5 | 0 | 3 | 20 | 0.1 | 0 | 0 | 0 | 1 | 5.2 | 5 | 0 | 3 | 20 |
| | Pull through for Hirschsprung's: scope | 3.6 | 3 | 0 | 4 | 10 | 0.1 | 0 | 0 | 0 | 1 | 3.7 | 3 | 0 | 4 | 10 |
| | Pull through for IBD or polyposis: open | 0.7 | 2 | 0 | 0 | 10 | 0.0 | 0 | 0 | 0 | 1 | 0.8 | 2 | 0 | 0 | 10 |
| | Pull through for IBD or polyposis: scope | 0.3 | 1 | 0 | 0 | 3 | 0.1 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 3 |
| | Exploratory lap with or without biopsy: open | 29.9 | 13 | 15 | 26 | 65 | 25.0 | 24 | 0 | 18 | 98 | 54.9 | 28 | 15 | 50 | 126 |
| | Exploratory lap with or without biopsy: lap | 3.5 | 3 | 0 | 3 | 10 | 1.8 | 3 | 0 | 1 | 12 | 5.3 | 5 | 0 | 4 | 19 |
| | Excision of omental/mesenteric cyst | 1.0 | 1 | 0 | 1 | 3 | 0.3 | 1 | 0 | 0 | 3 | 1.3 | 1 | 0 | 1 | 5 |
| | Omphalocele (any surgical repair) | 5.4 | 3 | 1 | 5 | 12 | 0.2 | 0 | 0 | 0 | 2 | 5.6 | 3 | 1 | 6 | 12 |
| | Gastroschisis (any surgical repair) | 13.2 | 9 | 0 | 12 | 39 | 0.1 | 0 | 0 | 0 | 2 | 13.3 | 9 | 0 | 12 | 39 |
| | Resection urachal remnant | 3.0 | 2 | 0 | 3 | 8 | 0.1 | 0 | 0 | 0 | 1 | 3.1 | 2 | 0 | 3 | 8 |
| | Resection omphalomesenteric duct/cyst | 2.5 | 2 | 0 | 2 | 11 | 0.3 | 1 | 0 | 0 | 3 | 2.8 | 3 | 0 | 2 | 11 |
| | Excision neuroblastoma/adrenal/other retroperitoneal tumor | 6.7 | 3 | 0 | 6 | 15 | 0.1 | 0 | 0 | 0 | 1 | 6.7 | 3 | 0 | 6 | 15 |
| | Excision sacrococcygeal teratoma | 2.7 | 2 | 0 | 3 | 8 | 0.0 | 0 | 0 | 0 | 0 | 2.7 | 2 | 0 | 3 | 8 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = All Roles

[PAGE 6]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|---------------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Abdominal | Diagnostic laparoscopy | 7.7 | 5 | 0 | 8 | 21 | 17.4 | 19 | 0 | 6 | 62 | 25.1 | 21 | 0 | 17 | 72 |
| | Adrenal: open | 0.7 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.8 | 1 | 0 | 0 | 4 |
| | Adrenal: scope | 1.2 | 1 | 0 | 1 | 4 | 0.1 | 0 | 0 | 0 | 1 | 1.3 | 1 | 0 | 1 | 4 |
| | Other (abdominal) | 20.3 | 10 | 4 | 19 | 42 | 2.1 | 2 | 0 | 1 | 12 | 22.3 | 10 | 7 | 20 | 43 |
| | TOTAL - Abdominal | 480.1 | 109 | 327 | 456 | 770 | 100.5 | 65 | 6 | 95 | 220 | 580.7 | 118 | 379 | 551 | 867 |
| Hernia Repair | Pediatric repair inguinal hernia > 6 months | 43.3 | 17 | 16 | 40 | 81 | 5.1 | 6 | 0 | 3 | 25 | 48.4 | 20 | 18 | 46 | 92 |
| | Infant repair inguinal hernia < 6 months | 41.1 | 20 | 19 | 35 | 96 | 7.4 | 9 | 0 | 4 | 37 | 48.4 | 22 | 22 | 41 | 97 |
| | Repair umbilical hernia | 20.0 | 13 | 1 | 17 | 62 | 4.0 | 5 | 0 | 3 | 18 | 24.1 | 15 | 1 | 21 | 78 |
| | Repair ventral hernia | 6.7 | 4 | 1 | 6 | 15 | 1.3 | 2 | 0 | 1 | 7 | 8.0 | 5 | 2 | 7 | 22 |
| | Repair femoral hernia | 0.2 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 2 |
| | Hernia repair: scope | 7.6 | 13 | 0 | 1 | 60 | 0.9 | 2 | 0 | 0 | 9 | 8.5 | 15 | 0 | 1 | 66 |
| | Other (hernia repair) | 1.8 | 2 | 0 | 1 | 8 | 0.5 | 1 | 0 | 0 | 3 | 2.3 | 2 | 0 | 2 | 11 |
| | TOTAL - Hernia Repair | 120.7 | 38 | 62 | 115 | 198 | 19.1 | 19 | 0 | 17 | 76 | 139.8 | 46 | 68 | 134 | 243 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = All Roles

[PAGE 7] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|---------------|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Liver/Biliary | Major hepatic resection/repair: tumor | 3.1 | 3 | 0 | 2 | 17 | 0.2 | 0 | 0 | 0 | 1 | 3.2 | 3 | 0 | 2 | 17 |
| | Major hepatic resection/repair: trauma | 0.6 | 1 | 0 | 0 | 3 | 0.1 | 0 | 0 | 0 | 1 | 0.7 | 1 | 0 | 0 | 3 |
| | Major hepatic resection/repair: other | 0.2 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 1 |
| | Lysis of adhesions | 0.5 | 1 | 0 | 0 | 3 | 1.7 | 4 | 0 | 0 | 26 | 2.2 | 5 | 0 | 1 | 29 |
| | Liver: billiary atresia | 3.3 | 3 | 0 | 3 | 10 | 0.3 | 0 | 0 | 0 | 2 | 3.5 | 3 | 0 | 3 | 10 |
| | Liver biopsy: open | 2.1 | 2 | 0 | 1 | 9 | 0.7 | 1 | 0 | 0 | 4 | 2.8 | 3 | 0 | 2 | 13 |
| | Liver harvest | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Liver transplant | 0.5 | 1 | 0 | 0 | 6 | 0.0 | 0 | 0 | 0 | 0 | 0.5 | 1 | 0 | 0 | 6 |
| | Cholecystectomy w/wo common bile duct exploration: open | 1.6 | 1 | 0 | 1 | 4 | 0.9 | 1 | 0 | 1 | 6 | 2.5 | 2 | 0 | 2 | 7 |
| | Cholecystectomy w/wo common bile duct exploration: scope | 26.5 | 12 | 9 | 23 | 58 | 1.4 | 2 | 0 | 1 | 6 | 27.9 | 13 | 9 | 24 | 58 |
| | Portoenterostomy | 0.8 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 2 | 0.8 | 1 | 0 | 0 | 4 |
| | Excision choledochal cyst | 2.4 | 2 | 0 | 2 | 7 | 0.7 | 1 | 0 | 1 | 2 | 3.0 | 2 | 0 | 2 | 8 |
| | Portosystemic shunts or other operations for portal hypertensior | 0.7 | 3 | 0 | 0 | 20 | 0.1 | 0 | 0 | 0 | 3 | 0.8 | 4 | 0 | 0 | 23 |
| | Pancreatic resection for: trauma | 0.9 | 1 | 0 | 1 | 3 | 0.1 | 0 | 0 | 0 | 1 | 0.9 | 1 | 0 | 1 | 3 |
| | Pancreatic resection for: hyperinsulinism | 0.5 | 3 | 0 | 0 | 18 | 0.0 | 0 | 0 | 0 | 0 | 0.5 | 3 | 0 | 0 | 18 |
| | Pancreatic resection for: tumor | 0.4 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 3 |
| | Pancreas: other | 0.3 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.3 | 1 | 0 | 0 | 2 |
| | Operations for pseudocyst | 0.3 | 1 | 0 | 0 | 4 | 0.0 | 0 | 0 | 0 | 0 | 0.3 | 1 | 0 | 0 | 4 |
| | Splenorrhaphy | 0.1 | 0 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 1 | 0.2 | 0 | 0 | 0 | 2 |
| | Splenectomy: open | 1.1 | 1 | 0 | 1 | 4 | 0.3 | 1 | 0 | 0 | 2 | 1.4 | 1 | 0 | 1 | 6 |
| | Splenectomy: scope | 5.7 | 3 | 1 | 6 | 12 | 0.1 | 0 | 0 | 0 | 1 | 5.8 | 3 | 1 | 6 | 12 |
| | Other (liver) | 0.3 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 2 |
| | TOTAL - Liver/Biliary | 51.7 | 15 | 25 | 51 | 85 | 6.6 | 6 | 0 | 6 | 34 | 58.3 | 17 | 28 | 59 | 96 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = All Roles

[PAGE 8]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|------------------------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Genito Urinary | Nephrectomy (total or partial): tumor | 5.6 | 3 | 0 | 6 | 12 | 0.4 | 1 | 0 | 0 | 6 | 6.0 | 3 | 0 | 6 | 12 |
| | Nephrectomy (total or partial): trauma | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Nephrectomy (total or partial/nephorrhaphy) | 0.5 | 1 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 2 | 0.6 | 1 | 0 | 0 | 4 |
| | Nephrectomy (total or partial): other | 0.5 | 1 | 0 | 0 | 5 | 0.1 | 0 | 0 | 0 | 2 | 0.6 | 1 | 0 | 0 | 5 |
| | Renal transplant | 2.0 | 5 | 0 | 0 | 25 | 0.1 | 0 | 0 | 0 | 2 | 2.1 | 5 | 0 | 0 | 25 |
| | Renal: scope | 0.5 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.7 | 1 | 0 | 0 | 4 |
| | Renal biopsy (open) | 0.3 | 1 | 0 | 0 | 2 | 0.2 | 0 | 0 | 0 | 2 | 0.5 | 1 | 0 | 0 | 2 |
| | Cystectomy (partial or total) | 0.3 | 1 | 0 | 0 | 3 | 0.1 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 3 |
| | Operation for nephro-uretero lithiasis | 0.1 | 0 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 1 | 0 | 0 | 4 |
| | Enteric conduit (any) | 0.4 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 2 | 0.4 | 1 | 0 | 0 | 4 |
| | Bladder augmentation or repair (any) | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 2 |
| | Pyeloplasty/UPJ reconstruction | 0.5 | 1 | 0 | 0 | 7 | 0.1 | 0 | 0 | 0 | 2 | 0.5 | 1 | 0 | 0 | 7 |
| | Ureteral reconstruction/reimplantation | 0.7 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.8 | 1 | 0 | 0 | 5 |
| | Reconstruct cloacal extrophy | 1.1 | 1 | 0 | 1 | 5 | 0.0 | 0 | 0 | 0 | 1 | 1.2 | 1 | 0 | 1 | 5 |
| | Circumcision (OR only) | 21.4 | 17 | 1 | 19 | 62 | 10.8 | 10 | 0 | 8 | 37 | 32.2 | 20 | 6 | 31 | 96 |
| | Orchidopexy: open | 16.4 | 11 | 4 | 14 | 49 | 1.5 | 3 | 0 | 1 | 20 | 17.9 | 13 | 4 | 16 | 69 |
| | Orchidopexy: scope | 2.5 | 3 | 0 | 2 | 11 | 0.2 | 1 | 0 | 0 | 4 | 2.7 | 3 | 0 | 2 | 15 |
| | Orchiectomy | 2.0 | 3 | 0 | 1 | 16 | 0.4 | 1 | 0 | 0 | 5 | 2.4 | 4 | 0 | 1 | 21 |
| | Operation for torsion testis/appendages | 1.0 | 2 | 0 | 0 | 11 | 0.2 | 1 | 0 | 0 | 3 | 1.2 | 3 | 0 | 0 | 11 |
| | Operation for varicocele: open | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Operation for varicocele: scope | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Procedures for intersex | 0.2 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 2 |
| | Vaginal procedures: open | 1.7 | 2 | 0 | 1 | 8 | 0.2 | 1 | 0 | 0 | 3 | 1.9 | 2 | 0 | 1 | 9 |
| | Oophorectomy (partial or total) | 11.4 | 7 | 1 | 10 | 35 | 0.7 | 1 | 0 | 0 | 4 | 12.2 | 7 | 2 | 11 | 35 |
| | Hysterectomy/salpingectomy: open | 0.4 | 1 | 0 | 0 | 2 | 0.2 | 0 | 0 | 0 | 1 | 0.5 | 1 | 0 | 0 | 3 |
| | Hysterectomy/salpingectomy: scope | 1.9 | 3 | 0 | 1 | 15 | 0.1 | 0 | 0 | 0 | 2 | 1.9 | 3 | 0 | 1 | 15 |
| | Rpr complex laceration vagina/perineum | 1.1 | 2 | 0 | 0 | 7 | 0.0 | 0 | 0 | 0 | 1 | 1.1 | 2 | 0 | 0 | 7 |
| Other (genito urinary) | 3.1 | 5 | 0 | 2 | 23 | 0.6 | 1 | 0 | 0 | 7 | 3.7 | 6 | 0 | 2 | 30 | |
| TOTAL - Genito Urinary | | 75.8 | 31 | 26 | 74 | 152 | 16.3 | 14 | 1 | 14 | 52 | 92.1 | 36 | 37 | 85 | 204 |

Based upon the number of procedures logged by residents in the indicated role.

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = All Roles

[PAGE 9] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|-----------------------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Endoscopic Procedures | Diagnostic thoracoscopy | 0.9 | 1 | 0 | 0 | 6 | 0.7 | 1 | 0 | 0 | 4 | 1.6 | 2 | 0 | 1 | 7 |
| | Esophageal dilation | 14.3 | 14 | 0 | 10 | 66 | 1.5 | 3 | 0 | 1 | 12 | 15.7 | 16 | 0 | 12 | 78 |
| | Esophagoscopy | 5.6 | 10 | 0 | 3 | 52 | 2.4 | 4 | 0 | 1 | 19 | 8.0 | 12 | 0 | 4 | 59 |
| | Esophagoscopy: gastroscopy | 5.7 | 9 | 0 | 2 | 44 | 1.6 | 3 | 0 | 0 | 14 | 7.3 | 11 | 0 | 3 | 57 |
| | Cystoscopy/colposcopy | 3.6 | 5 | 0 | 1 | 23 | 1.1 | 2 | 0 | 0 | 8 | 4.6 | 6 | 0 | 3 | 24 |
| | Laryngoscopy/bronchoscopy | 8.3 | 7 | 1 | 6 | 31 | 3.5 | 4 | 0 | 3 | 14 | 11.8 | 9 | 1 | 9 | 37 |
| | Removal foreign body esophagus or trachea | 21.5 | 16 | 0 | 20 | 61 | 0.8 | 1 | 0 | 0 | 3 | 22.3 | 16 | 0 | 21 | 62 |
| | Colonoscopy | 0.6 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.6 | 1 | 0 | 0 | 4 |
| | Sigmoidoscopy | 3.0 | 4 | 0 | 2 | 16 | 0.9 | 2 | 0 | 0 | 8 | 3.9 | 5 | 0 | 3 | 22 |
| | TOTAL - Endoscopic Procedures | 63.4 | 32 | 21 | 53 | 148 | 12.5 | 12 | 0 | 10 | 65 | 75.9 | 40 | 25 | 71 | 213 |
| Trauma | Non-operative Trauma | 128.5 | 68 | 8 | 113 | 300 | 0.2 | 1 | 0 | 0 | 4 | 128.7 | 69 | 8 | 113 | 304 |
| Total Procedures | TOTAL - Tracked Codes | 1,177.6 | 212 | 930 | 1,100 | 1,715 | 187.6 | 118 | 12 | 173 | 410 | 1,365.2 | 252 | 1,007 | 1,313 | 2,036 |
| Non-Tracked | TOTAL - Non-Tracked Codes | 54.6 | 39 | 0 | 53 | 155 | 32.1 | 37 | 0 | 14 | 168 | 86.7 | 63 | 4 | 81 | 233 |

Based upon the number of procedures logged by residents in the indicated role.

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Surgeon

[PAGE 1] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|----------------------------------|--------------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Skin/Soft Tissue/Musculoskeletal | Burn debridement or grafting | 9.5 | 22 | 0 | 1 | 115 | 3.7 | 16 | 0 | 0 | 98 | 13.1 | 37 | 0 | 1 | 213 |
| | Complex wound closure | 4.3 | 6 | 0 | 3 | 30 | 1.0 | 2 | 0 | 0 | 8 | 5.3 | 6 | 0 | 4 | 31 |
| | Breast | 3.3 | 3 | 0 | 3 | 15 | 0.1 | 0 | 0 | 0 | 1 | 3.4 | 3 | 0 | 3 | 15 |
| | Pilonidal cyst excision | 3.5 | 4 | 0 | 3 | 20 | 0.0 | 0 | 0 | 0 | 0 | 3.5 | 4 | 0 | 3 | 20 |
| | Major excision soft tissue tumor | 8.2 | 4 | 1 | 7 | 22 | 0.4 | 1 | 0 | 0 | 2 | 8.5 | 4 | 1 | 8 | 22 |
| | Major soft tissue repair for trauma | 4.6 | 6 | 0 | 2 | 23 | 0.6 | 2 | 0 | 0 | 8 | 5.3 | 7 | 0 | 2 | 30 |
| | Other (skin/soft tissue/musc) | 10.8 | 7 | 0 | 9 | 32 | 1.1 | 1 | 0 | 1 | 5 | 11.8 | 8 | 0 | 11 | 36 |
| | TOTAL - Skin/Soft Tissue/Musc | 44.1 | 33 | 10 | 37 | 175 | 6.8 | 19 | 0 | 2 | 115 | 50.9 | 49 | 10 | 38 | 290 |
| Head & Neck | Thyroidectomy (any) | 5.8 | 7 | 0 | 3 | 23 | 0.1 | 0 | 0 | 0 | 1 | 5.9 | 7 | 0 | 3 | 23 |
| | Parathyroidectomy (any) | 0.6 | 1 | 0 | 0 | 4 | 0.0 | 0 | 0 | 0 | 0 | 0.6 | 1 | 0 | 0 | 4 |
| | Branchial cleft cyst/sinus | 3.6 | 2 | 0 | 4 | 7 | 0.1 | 0 | 0 | 0 | 1 | 3.6 | 2 | 0 | 4 | 7 |
| | Thyroglossal duct cyst/sinus | 2.5 | 2 | 0 | 2 | 14 | 0.0 | 0 | 0 | 0 | 0 | 2.5 | 2 | 0 | 2 | 14 |
| | Cystic hygroma/lymphangioma | 0.7 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 1 | 0.8 | 1 | 0 | 0 | 3 |
| | Dermoid/other cyst | 7.4 | 7 | 0 | 5 | 31 | 0.6 | 2 | 0 | 0 | 8 | 8.0 | 7 | 0 | 5 | 31 |
| | Major tumor (head & neck) | 1.0 | 1 | 0 | 1 | 4 | 0.1 | 0 | 0 | 0 | 1 | 1.1 | 1 | 0 | 1 | 4 |
| | Other (head & neck) | 3.9 | 3 | 0 | 3 | 11 | 0.2 | 0 | 0 | 0 | 1 | 4.1 | 3 | 0 | 3 | 11 |
| | TOTAL - Head & Neck | 25.7 | 11 | 11 | 24 | 55 | 0.9 | 2 | 0 | 0 | 8 | 26.6 | 11 | 11 | 24 | 57 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Surgeon

| [PAGE 2] | | Number of Programs in the Nation: 34 | | | | | Number of Residents in the Nation: 39 | | | | | | | | | |
|-----------|---|--------------------------------------|--------------|--------------|--------------|--------------|---------------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Primary | | | | | Secondary | | | | | Total | | | | |
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Thoracic | Tracheostomy | 1.9 | 3 | 0 | 0 | 11 | 0.3 | 1 | 0 | 0 | 2 | 2.2 | 3 | 0 | 1 | 12 |
| | Laryngeal or tracheal resect and/or recon | 0.2 | 1 | 0 | 0 | 2 | 0.1 | 1 | 0 | 0 | 4 | 0.3 | 1 | 0 | 0 | 6 |
| | Repair chest wall deformity | 13.7 | 9 | 2 | 13 | 49 | 0.1 | 0 | 0 | 0 | 2 | 13.8 | 9 | 2 | 13 | 51 |
| | Resection chest wall tumor | 1.5 | 1 | 0 | 1 | 5 | 0.1 | 0 | 0 | 0 | 2 | 1.6 | 1 | 0 | 1 | 5 |
| | Excision mediastinal cyst | 2.9 | 2 | 0 | 2 | 9 | 0.1 | 0 | 0 | 0 | 2 | 3.0 | 2 | 0 | 3 | 10 |
| | Excision mediastinal tumor | 3.2 | 2 | 0 | 3 | 10 | 0.4 | 1 | 0 | 0 | 2 | 3.5 | 3 | 0 | 4 | 10 |
| | Pulmonary resection: open | 8.0 | 7 | 1 | 7 | 40 | 0.7 | 1 | 0 | 0 | 8 | 8.7 | 7 | 1 | 7 | 40 |
| | Pulmonary resection: scope | 7.3 | 4 | 1 | 6 | 18 | 0.2 | 0 | 0 | 0 | 1 | 7.5 | 4 | 1 | 6 | 19 |
| | Thoracotomy for trauma | 0.7 | 1 | 0 | 1 | 3 | 0.1 | 0 | 0 | 0 | 2 | 0.8 | 1 | 0 | 1 | 3 |
| | Lung biopsy: open | 2.0 | 2 | 0 | 2 | 7 | 0.8 | 2 | 0 | 0 | 10 | 2.8 | 3 | 0 | 2 | 14 |
| | Lung biopsy: scope | 3.9 | 4 | 0 | 2 | 14 | 0.4 | 1 | 0 | 0 | 4 | 4.4 | 4 | 0 | 3 | 17 |
| | Decortication/pleurectomy/blebectomy: open | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 2 |
| | Decortication/pleurectomy/blebectomy: scope | 5.1 | 3 | 0 | 4 | 15 | 0.5 | 1 | 0 | 0 | 4 | 5.6 | 4 | 0 | 5 | 15 |
| | Esophageal resection or replacement | 2.6 | 2 | 0 | 2 | 10 | 0.2 | 0 | 0 | 0 | 2 | 2.8 | 2 | 0 | 2 | 11 |
| | Esophagomyotomy | 1.3 | 1 | 0 | 1 | 4 | 0.1 | 0 | 0 | 0 | 1 | 1.4 | 1 | 0 | 1 | 4 |
| | Repair esophageal atresia/tracheoesophageal fistula | 9.9 | 6 | 2 | 9 | 33 | 0.1 | 0 | 0 | 0 | 1 | 10.1 | 6 | 2 | 9 | 34 |
| | Thoracoscopy: other | 0.6 | 1 | 0 | 0 | 4 | 0.5 | 2 | 0 | 0 | 9 | 1.1 | 2 | 0 | 0 | 11 |
| | Thoracotomy: other | 1.3 | 2 | 0 | 1 | 11 | 0.5 | 1 | 0 | 0 | 8 | 1.8 | 3 | 0 | 1 | 13 |
| | Other (thoracic) | 1.1 | 3 | 0 | 0 | 14 | 0.4 | 1 | 0 | 0 | 5 | 1.5 | 3 | 0 | 0 | 14 |
| | TOTAL - Thoracic | 67.4 | 21 | 24 | 64 | 115 | 5.7 | 7 | 0 | 3 | 31 | 73.1 | 24 | 27 | 70 | 132 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Surgeon

[PAGE 3] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|----------------|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Diaphragm | Repair diaphragmatic hernia | 12.7 | 6 | 4 | 12 | 31 | 0.2 | 1 | 0 | 0 | 3 | 12.9 | 6 | 4 | 12 | 33 |
| | Plication of diaphragm | 0.7 | 1 | 0 | 1 | 2 | 0.1 | 0 | 0 | 0 | 1 | 0.8 | 1 | 0 | 1 | 3 |
| | Transthoracic and/or retroperitoneal exp for scoliosis | 0.8 | 2 | 0 | 0 | 6 | 0.1 | 0 | 0 | 0 | 2 | 0.9 | 2 | 0 | 0 | 7 |
| | Other (diaphragm) | 0.2 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 4 | 0.5 | 1 | 0 | 0 | 4 |
| | TOTAL - Diaphragm | 14.5 | 7 | 5 | 14 | 33 | 0.7 | 1 | 0 | 0 | 4 | 15.2 | 7 | 5 | 15 | 36 |
| Cardiovascular | Patent ductus arteriosus | 2.0 | 5 | 0 | 0 | 30 | 0.0 | 0 | 0 | 0 | 1 | 2.0 | 5 | 0 | 0 | 30 |
| | Vascular ring | 0.3 | 1 | 0 | 0 | 5 | 0.1 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 6 |
| | Any open heart procedure | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Any closed heart procedure | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 1 |
| | Aortopexy | 0.7 | 1 | 0 | 0 | 7 | 0.0 | 0 | 0 | 0 | 0 | 0.7 | 1 | 0 | 0 | 7 |
| | Renal artery reconstruction | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Peripheral artery reconstruction | 0.2 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 2 |
| | Construction or take down AV fistula/shunt | 0.5 | 1 | 0 | 0 | 5 | 0.0 | 0 | 0 | 0 | 0 | 0.5 | 1 | 0 | 0 | 5 |
| | Surgical placement/removal central access line | 76.1 | 34 | 32 | 75 | 170 | 16.5 | 17 | 0 | 14 | 76 | 92.6 | 43 | 34 | 91 | 246 |
| | Dialysis access insertion/removal | 1.2 | 2 | 0 | 1 | 10 | 0.1 | 0 | 0 | 0 | 2 | 1.3 | 2 | 0 | 1 | 10 |
| | Cannulate/ECMO | 15.4 | 8 | 0 | 15 | 36 | 0.2 | 1 | 0 | 0 | 2 | 15.6 | 8 | 0 | 15 | 36 |
| | Major vessel reconstruction | 1.3 | 2 | 0 | 0 | 13 | 0.2 | 1 | 0 | 0 | 3 | 1.5 | 3 | 0 | 1 | 16 |
| | Vascular trauma | 0.6 | 1 | 0 | 0 | 8 | 0.0 | 0 | 0 | 0 | 0 | 0.6 | 1 | 0 | 0 | 8 |
| | Vascular: post-operative exploration | 0.2 | 1 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 2 |
| | TOTAL - Cardiovascular | 98.6 | 37 | 47 | 95 | 217 | 17.3 | 17 | 0 | 14 | 79 | 115.8 | 47 | 48 | 109 | 296 |

Based upon the number of procedures logged by residents in the indicated role.

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Surgeon

[PAGE 4] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|-----------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Abdominal | Antireflux procedure: open | 7.4 | 9 | 0 | 5 | 42 | 0.9 | 1 | 0 | 1 | 4 | 8.3 | 9 | 0 | 5 | 43 |
| | Antireflux procedure: scope | 26.2 | 21 | 2 | 18 | 96 | 0.9 | 1 | 0 | 0 | 5 | 27.1 | 22 | 2 | 18 | 96 |
| | Pyloroplasty/gastric resection w/wout vagotomy | 2.4 | 2 | 0 | 2 | 9 | 0.6 | 1 | 0 | 0 | 3 | 2.9 | 3 | 0 | 2 | 10 |
| | Pyloroplasty/gastric resect w/wout vagotomy - lap | 1.7 | 5 | 0 | 0 | 29 | 0.1 | 0 | 0 | 0 | 2 | 1.8 | 5 | 0 | 0 | 29 |
| | Any gastrostomy/jejunostomy: open | 13.1 | 10 | 3 | 11 | 45 | 9.8 | 9 | 0 | 8 | 35 | 22.9 | 15 | 4 | 18 | 68 |
| | Any gastrostomy/jejunostomy: scope | 34.1 | 23 | 6 | 24 | 99 | 14.8 | 15 | 0 | 10 | 58 | 48.9 | 30 | 11 | 43 | 134 |
| | Pyloromyotomy: open | 7.9 | 11 | 0 | 3 | 43 | 0.1 | 0 | 0 | 0 | 1 | 7.9 | 11 | 0 | 3 | 43 |
| | Pyloromyotomy: scope | 30.6 | 21 | 0 | 25 | 107 | 0.2 | 0 | 0 | 0 | 1 | 30.7 | 21 | 0 | 25 | 107 |
| | Operation for malrotation | 10.2 | 4 | 1 | 11 | 20 | 1.4 | 2 | 0 | 1 | 9 | 11.6 | 5 | 1 | 12 | 22 |
| | Duodenal atresia | 5.3 | 3 | 0 | 5 | 13 | 0.1 | 0 | 0 | 0 | 2 | 5.4 | 3 | 0 | 6 | 13 |
| | Repair intestinal atresia, stenosis or web | 11.5 | 8 | 1 | 10 | 41 | 1.9 | 2 | 0 | 2 | 6 | 13.4 | 8 | 1 | 12 | 42 |
| | Intestinal resection: open | 23.1 | 11 | 6 | 21 | 52 | 5.7 | 6 | 0 | 4 | 26 | 28.8 | 13 | 9 | 28 | 53 |
| | Intestinal resection: scope | 2.5 | 2 | 0 | 2 | 8 | 0.1 | 0 | 0 | 0 | 1 | 2.6 | 2 | 0 | 2 | 8 |
| | Intestinal resection/repair or ostomy for IBD | 2.4 | 2 | 0 | 2 | 9 | 0.3 | 1 | 0 | 0 | 3 | 2.8 | 2 | 0 | 3 | 11 |
| | Intestinal resect/repair or ostomy for IBD - lap | 5.3 | 4 | 0 | 5 | 16 | 0.1 | 0 | 0 | 0 | 1 | 5.4 | 4 | 0 | 5 | 16 |
| | Intestinal resection/repair or ostomy for trauma | 4.4 | 3 | 0 | 4 | 16 | 1.9 | 2 | 0 | 1 | 8 | 6.3 | 5 | 0 | 5 | 18 |
| | Intestinal resect/repair or ostomy for trauma - lap | 0.4 | 1 | 0 | 0 | 4 | 0.0 | 0 | 0 | 0 | 1 | 0.5 | 1 | 0 | 0 | 4 |
| | Bowel resection: open | 4.4 | 5 | 0 | 3 | 28 | 0.7 | 1 | 0 | 0 | 5 | 5.1 | 5 | 0 | 4 | 28 |
| | Bowel resection: scope | 1.7 | 2 | 0 | 1 | 9 | 0.1 | 0 | 0 | 0 | 1 | 1.8 | 2 | 0 | 1 | 9 |
| | Lap/resec for intussusception-malrotation-volvulus | 7.1 | 3 | 2 | 6 | 14 | 0.4 | 1 | 0 | 0 | 4 | 7.5 | 3 | 2 | 7 | 15 |
| | Ostomy for anorectal malformation: open | 3.8 | 2 | 0 | 4 | 10 | 0.5 | 1 | 0 | 0 | 3 | 4.3 | 3 | 1 | 4 | 12 |
| | Ostomy for anorectal malformation: laparoscopic | 0.6 | 1 | 0 | 0 | 6 | 0.2 | 0 | 0 | 0 | 1 | 0.8 | 1 | 0 | 0 | 6 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Surgeon

[PAGE 5]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|-----------|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Abdominal | Ostomy for Hirschsprung's: open | 1.8 | 2 | 0 | 1 | 6 | 0.1 | 0 | 0 | 0 | 1 | 1.8 | 2 | 0 | 1 | 6 |
| | Ostomy for Hirschsprung's: laparoscopic | 1.3 | 1 | 0 | 1 | 8 | 0.1 | 0 | 0 | 0 | 2 | 1.4 | 2 | 0 | 1 | 8 |
| | Ostomy for other: open | 6.8 | 4 | 1 | 7 | 16 | 1.8 | 2 | 0 | 1 | 7 | 8.6 | 5 | 1 | 9 | 19 |
| | Ostomy for other: laparoscopic | 0.6 | 1 | 0 | 0 | 3 | 0.2 | 0 | 0 | 0 | 2 | 0.7 | 1 | 0 | 0 | 4 |
| | Closure/revision any ostomy/fistula: open | 24.6 | 9 | 6 | 25 | 51 | 2.8 | 3 | 0 | 2 | 11 | 27.5 | 10 | 6 | 27 | 58 |
| | Closure/revision any ostomy/fistula: scope | 0.4 | 1 | 0 | 0 | 7 | 0.0 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 8 |
| | Appendectomy: open | 2.8 | 4 | 0 | 1 | 17 | 3.8 | 6 | 0 | 1 | 26 | 6.6 | 8 | 0 | 3 | 31 |
| | Appendectomy: scope | 93.5 | 59 | 13 | 81 | 288 | 2.3 | 3 | 0 | 1 | 15 | 95.8 | 60 | 14 | 83 | 295 |
| | Perineal procedure for imperforate anus | 13.7 | 6 | 3 | 14 | 30 | 0.2 | 0 | 0 | 0 | 2 | 13.9 | 6 | 3 | 14 | 31 |
| | Pull through for Hirschsprung's: open | 5.0 | 4 | 0 | 3 | 19 | 0.1 | 0 | 0 | 0 | 1 | 5.1 | 4 | 0 | 3 | 19 |
| | Pull through for Hirschsprung's: scope | 3.5 | 2 | 0 | 4 | 10 | 0.1 | 0 | 0 | 0 | 1 | 3.6 | 2 | 0 | 4 | 10 |
| | Pull through for IBD or polyposis: open | 0.7 | 2 | 0 | 0 | 10 | 0.0 | 0 | 0 | 0 | 1 | 0.8 | 2 | 0 | 0 | 10 |
| | Pull through for IBD or polyposis: scope | 0.3 | 1 | 0 | 0 | 3 | 0.1 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 3 |
| | Exploratory lap with or without biopsy: open | 29.0 | 12 | 15 | 26 | 62 | 24.5 | 23 | 0 | 17 | 95 | 53.5 | 27 | 15 | 49 | 121 |
| | Exploratory lap with or without biopsy: lap | 3.3 | 3 | 0 | 3 | 9 | 1.8 | 3 | 0 | 1 | 12 | 5.1 | 4 | 0 | 4 | 19 |
| | Excision of omental/mesenteric cyst | 0.9 | 1 | 0 | 1 | 3 | 0.3 | 1 | 0 | 0 | 2 | 1.2 | 1 | 0 | 1 | 5 |
| | Omphalocele (any surgical repair) | 5.3 | 3 | 1 | 5 | 12 | 0.2 | 0 | 0 | 0 | 2 | 5.5 | 3 | 1 | 6 | 12 |
| | Gastroschisis (any surgical repair) | 12.7 | 8 | 0 | 11 | 39 | 0.1 | 0 | 0 | 0 | 2 | 12.9 | 8 | 0 | 11 | 39 |
| | Resection urachal remnant | 2.9 | 2 | 0 | 3 | 8 | 0.1 | 0 | 0 | 0 | 1 | 3.0 | 2 | 0 | 3 | 8 |
| | Resection omphalomesenteric duct/cyst | 2.4 | 2 | 0 | 2 | 11 | 0.3 | 1 | 0 | 0 | 3 | 2.7 | 2 | 0 | 2 | 11 |
| | Excision neuroblastoma/adrenal/other retroperitoneal tumor | 6.5 | 3 | 0 | 6 | 15 | 0.1 | 0 | 0 | 0 | 1 | 6.6 | 3 | 0 | 6 | 15 |
| | Excision sacrococcygeal teratoma | 2.7 | 2 | 0 | 3 | 7 | 0.0 | 0 | 0 | 0 | 0 | 2.7 | 2 | 0 | 3 | 7 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Surgeon

[PAGE 6]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|---------------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Abdominal | Diagnostic laparoscopy | 7.4 | 5 | 0 | 7 | 20 | 16.7 | 18 | 0 | 6 | 58 | 24.2 | 20 | 0 | 17 | 68 |
| | Adrenal: open | 0.7 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.8 | 1 | 0 | 0 | 4 |
| | Adrenal: scope | 1.2 | 1 | 0 | 1 | 4 | 0.1 | 0 | 0 | 0 | 1 | 1.2 | 1 | 0 | 1 | 4 |
| | Other (abdominal) | 19.6 | 9 | 4 | 19 | 42 | 2.0 | 2 | 0 | 1 | 12 | 21.6 | 10 | 7 | 20 | 43 |
| | TOTAL - Abdominal | 455.8 | 105 | 312 | 436 | 761 | 98.5 | 63 | 6 | 95 | 203 | 554.3 | 114 | 375 | 537 | 858 |
| Hernia Repair | Pediatric repair inguinal hernia > 6 months | 42.4 | 16 | 16 | 39 | 81 | 5.0 | 6 | 0 | 3 | 25 | 47.4 | 19 | 18 | 46 | 86 |
| | Infant repair inguinal hernia < 6 months | 40.6 | 19 | 19 | 35 | 96 | 7.2 | 9 | 0 | 4 | 35 | 47.7 | 21 | 22 | 41 | 97 |
| | Repair umbilical hernia | 19.1 | 12 | 1 | 17 | 53 | 4.0 | 4 | 0 | 3 | 18 | 23.1 | 14 | 1 | 21 | 68 |
| | Repair ventral hernia | 6.5 | 4 | 1 | 6 | 15 | 1.2 | 2 | 0 | 1 | 7 | 7.7 | 5 | 2 | 7 | 20 |
| | Repair femoral hernia | 0.2 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 1 |
| | Hernia repair: scope | 7.6 | 13 | 0 | 1 | 60 | 0.8 | 2 | 0 | 0 | 9 | 8.4 | 15 | 0 | 1 | 66 |
| | Other (hernia repair) | 1.8 | 2 | 0 | 1 | 8 | 0.5 | 1 | 0 | 0 | 3 | 2.3 | 2 | 0 | 2 | 11 |
| | TOTAL - Hernia Repair | 118.1 | 36 | 62 | 114 | 197 | 18.7 | 18 | 0 | 17 | 72 | 136.9 | 43 | 68 | 133 | 234 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Surgeon

[PAGE 7]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|---------------|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Liver/Biliary | Major hepatic resection/repair: tumor | 3.0 | 3 | 0 | 2 | 16 | 0.2 | 0 | 0 | 0 | 1 | 3.2 | 3 | 0 | 2 | 16 |
| | Major hepatic resection/repair: trauma | 0.6 | 1 | 0 | 0 | 3 | 0.1 | 0 | 0 | 0 | 1 | 0.7 | 1 | 0 | 0 | 3 |
| | Major hepatic resection/repair: other | 0.2 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 1 |
| | Lysis of adhesions | 0.4 | 1 | 0 | 0 | 2 | 1.6 | 4 | 0 | 0 | 24 | 2.1 | 4 | 0 | 1 | 26 |
| | Liver: billiary atresia | 3.2 | 3 | 0 | 3 | 10 | 0.3 | 0 | 0 | 0 | 2 | 3.5 | 3 | 0 | 3 | 10 |
| | Liver biopsy: open | 2.0 | 2 | 0 | 1 | 9 | 0.7 | 1 | 0 | 0 | 4 | 2.7 | 3 | 0 | 2 | 13 |
| | Liver harvest | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Liver transplant | 0.5 | 1 | 0 | 0 | 6 | 0.0 | 0 | 0 | 0 | 0 | 0.5 | 1 | 0 | 0 | 6 |
| | Cholecystectomy w/wo common bile duct exploration: open | 1.5 | 1 | 0 | 1 | 4 | 0.9 | 1 | 0 | 1 | 6 | 2.5 | 2 | 0 | 2 | 7 |
| | Cholecystectomy w/wo common bile duct exploration: scope | 24.7 | 12 | 9 | 22 | 58 | 1.3 | 2 | 0 | 1 | 5 | 26.1 | 12 | 9 | 23 | 58 |
| | Portoenterostomy | 0.8 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 2 | 0.8 | 1 | 0 | 0 | 4 |
| | Excision choledochal cyst | 2.3 | 2 | 0 | 2 | 7 | 0.6 | 1 | 0 | 1 | 2 | 2.9 | 2 | 0 | 2 | 8 |
| | Portosystemic shunts or other operations for portal hypertensior | 0.7 | 3 | 0 | 0 | 20 | 0.1 | 0 | 0 | 0 | 3 | 0.8 | 4 | 0 | 0 | 23 |
| | Pancreatic resection for: trauma | 0.8 | 1 | 0 | 1 | 3 | 0.1 | 0 | 0 | 0 | 1 | 0.9 | 1 | 0 | 1 | 3 |
| | Pancreatic resection for: hyperinsulinism | 0.5 | 3 | 0 | 0 | 18 | 0.0 | 0 | 0 | 0 | 0 | 0.5 | 3 | 0 | 0 | 18 |
| | Pancreatic resection for: tumor | 0.4 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 3 |
| | Pancreas: other | 0.3 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.3 | 1 | 0 | 0 | 2 |
| | Operations for pseudocyst | 0.3 | 1 | 0 | 0 | 4 | 0.0 | 0 | 0 | 0 | 0 | 0.3 | 1 | 0 | 0 | 4 |
| | Splenorrhaphy | 0.1 | 0 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 1 | 0.2 | 0 | 0 | 0 | 2 |
| | Splenectomy: open | 1.1 | 1 | 0 | 1 | 4 | 0.3 | 1 | 0 | 0 | 2 | 1.4 | 1 | 0 | 1 | 6 |
| | Splenectomy: scope | 5.6 | 3 | 1 | 6 | 12 | 0.1 | 0 | 0 | 0 | 1 | 5.7 | 3 | 1 | 6 | 12 |
| | Other (liver) | 0.3 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 2 |
| | TOTAL - Liver/Biliary | 49.4 | 14 | 20 | 49 | 78 | 6.4 | 6 | 0 | 5 | 32 | 55.8 | 16 | 26 | 55 | 92 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Surgeon

[PAGE 8]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|--|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Genito Urinary | Nephrectomy (total or partial): tumor | 5.4 | 3 | 0 | 6 | 12 | 0.4 | 1 | 0 | 0 | 6 | 5.7 | 3 | 0 | 6 | 12 |
| | Nephrectomy (total or partial): trauma | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Nephrectomy (total or partial/nephorrhaphy) | 0.4 | 1 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 2 | 0.6 | 1 | 0 | 0 | 4 |
| | Nephrectomy (total or partial): other | 0.5 | 1 | 0 | 0 | 5 | 0.1 | 0 | 0 | 0 | 2 | 0.6 | 1 | 0 | 0 | 5 |
| | Renal transplant | 1.9 | 5 | 0 | 0 | 23 | 0.1 | 0 | 0 | 0 | 2 | 2.0 | 5 | 0 | 0 | 23 |
| | Renal: scope | 0.5 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.7 | 1 | 0 | 0 | 4 |
| | Renal biopsy (open) | 0.3 | 1 | 0 | 0 | 2 | 0.2 | 0 | 0 | 0 | 2 | 0.5 | 1 | 0 | 0 | 2 |
| | Cystectomy (partial or total) | 0.3 | 1 | 0 | 0 | 3 | 0.1 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 3 |
| | Operation for nephro-uretero lithiasis | 0.1 | 0 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 1 | 0 | 0 | 4 |
| | Enteric conduit (any) | 0.3 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 2 | 0.4 | 1 | 0 | 0 | 4 |
| | Bladder augmentation or repair (any) | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 2 |
| | Pyeloplasty/UPJ reconstruction | 0.5 | 1 | 0 | 0 | 7 | 0.1 | 0 | 0 | 0 | 2 | 0.5 | 1 | 0 | 0 | 7 |
| | Ureteral reconstruction/reimplantation | 0.7 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.8 | 1 | 0 | 0 | 5 |
| | Reconstruct cloacal extrophy | 1.1 | 1 | 0 | 1 | 5 | 0.0 | 0 | 0 | 0 | 1 | 1.1 | 1 | 0 | 1 | 5 |
| | Circumcision (OR only) | 20.9 | 17 | 1 | 19 | 61 | 10.8 | 10 | 0 | 8 | 37 | 31.7 | 20 | 5 | 31 | 96 |
| | Orchidopexy: open | 16.3 | 10 | 4 | 14 | 47 | 1.5 | 3 | 0 | 1 | 18 | 17.7 | 12 | 4 | 16 | 65 |
| | Orchidopexy: scope | 2.5 | 3 | 0 | 2 | 11 | 0.2 | 1 | 0 | 0 | 4 | 2.7 | 3 | 0 | 2 | 15 |
| | Orchiectomy | 2.0 | 3 | 0 | 1 | 16 | 0.4 | 1 | 0 | 0 | 5 | 2.4 | 4 | 0 | 1 | 21 |
| | Operation for torsion testis/appendages | 1.0 | 2 | 0 | 0 | 11 | 0.2 | 1 | 0 | 0 | 3 | 1.2 | 3 | 0 | 0 | 11 |
| | Operation for varicocele: open | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Operation for varicocele: scope | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Procedures for intersex | 0.2 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 2 |
| | Vaginal procedures: open | 1.7 | 2 | 0 | 1 | 8 | 0.2 | 1 | 0 | 0 | 3 | 1.8 | 2 | 0 | 1 | 9 |
| Oophorectomy (partial or total) | 10.8 | 7 | 1 | 9 | 34 | 0.7 | 1 | 0 | 0 | 4 | 11.5 | 7 | 2 | 10 | 34 | |
| Hysterectomy/salpingectomy: open | 0.4 | 1 | 0 | 0 | 2 | 0.2 | 0 | 0 | 0 | 1 | 0.5 | 1 | 0 | 0 | 3 | |
| Hysterectomy/salpingectomy: scope | 1.8 | 3 | 0 | 1 | 13 | 0.1 | 0 | 0 | 0 | 2 | 1.8 | 3 | 0 | 1 | 13 | |
| Rpr complex laceration vagina/perineum | 1.0 | 2 | 0 | 0 | 7 | 0.0 | 0 | 0 | 0 | 1 | 1.1 | 2 | 0 | 0 | 7 | |
| Other (genito urinary) | 3.1 | 5 | 0 | 2 | 23 | 0.6 | 1 | 0 | 0 | 7 | 3.7 | 6 | 0 | 2 | 30 | |
| TOTAL - Genito Urinary | | 73.9 | 30 | 26 | 66 | 152 | 16.2 | 14 | 1 | 14 | 52 | 90.1 | 36 | 37 | 84 | 204 |

Based upon the number of procedures logged by residents in the indicated role.

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Surgeon

[PAGE 9] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|-----------------------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Endoscopic Procedures | Diagnostic thoracoscopy | 0.9 | 1 | 0 | 0 | 6 | 0.7 | 1 | 0 | 0 | 4 | 1.6 | 2 | 0 | 1 | 7 |
| | Esophageal dilation | 14.2 | 14 | 0 | 10 | 65 | 1.4 | 3 | 0 | 1 | 11 | 15.6 | 16 | 0 | 12 | 76 |
| | Esophagoscopy | 5.4 | 9 | 0 | 3 | 50 | 2.3 | 4 | 0 | 1 | 18 | 7.7 | 11 | 0 | 4 | 56 |
| | Esophagoscopy: gastroscopy | 5.6 | 9 | 0 | 2 | 44 | 1.6 | 3 | 0 | 0 | 14 | 7.2 | 11 | 0 | 3 | 57 |
| | Cystoscopy/colposcopy | 3.5 | 5 | 0 | 1 | 23 | 1.1 | 2 | 0 | 0 | 7 | 4.6 | 6 | 0 | 3 | 24 |
| | Laryngoscopy/bronchoscopy | 8.3 | 7 | 1 | 6 | 31 | 3.4 | 4 | 0 | 3 | 13 | 11.7 | 8 | 1 | 9 | 37 |
| | Removal foreign body esophagus or trachea | 21.0 | 16 | 0 | 20 | 61 | 0.8 | 1 | 0 | 0 | 3 | 21.8 | 16 | 0 | 21 | 62 |
| | Colonoscopy | 0.6 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.6 | 1 | 0 | 0 | 4 |
| | Sigmoidoscopy | 2.9 | 4 | 0 | 2 | 14 | 0.9 | 2 | 0 | 0 | 8 | 3.8 | 5 | 0 | 3 | 22 |
| | TOTAL - Endoscopic Procedures | 62.4 | 31 | 21 | 52 | 142 | 12.3 | 12 | 0 | 10 | 61 | 74.7 | 39 | 25 | 68 | 203 |
| Trauma | Non-operative Trauma | 126.9 | 69 | 8 | 110 | 299 | 0.2 | 1 | 0 | 0 | 4 | 127.2 | 69 | 8 | 110 | 303 |
| Total Procedures | TOTAL - Tracked Codes | 1,136.8 | 202 | 866 | 1,059 | 1,649 | 183.8 | 115 | 12 | 173 | 394 | 1,320.6 | 239 | 934 | 1,254 | 1,943 |
| Non-Tracked | TOTAL - Non-Tracked Codes | 52.0 | 37 | 0 | 50 | 154 | 31.5 | 36 | 0 | 14 | 164 | 83.5 | 60 | 4 | 81 | 230 |

Based upon the number of procedures logged by residents in the indicated role.

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Teaching Assistant

[PAGE 1] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|----------------------------------|-------------------------------------|---------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Skin/Soft Tissue/Musculoskeletal | Burn debridement or grafting | 0.4 | 1 | 0 | 0 | 6 | 0.3 | 1 | 0 | 0 | 8 | 0.7 | 2 | 0 | 0 | 14 |
| | Complex wound closure | 0.4 | 1 | 0 | 0 | 5 | 0.1 | 0 | 0 | 0 | 2 | 0.5 | 1 | 0 | 0 | 5 |
| | Breast | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Pilonidal cyst excision | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Major excision soft tissue tumor | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Major soft tissue repair for trauma | 0.2 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 1 | 0 | 0 | 2 |
| | Other (skin/soft tissue/musc) | 0.2 | 1 | 0 | 0 | 4 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 1 | 0 | 0 | 4 |
| | TOTAL - Skin/Soft Tissue/Musc | 1.5 | 3 | 0 | 1 | 15 | 0.4 | 2 | 0 | 0 | 9 | 1.9 | 4 | 0 | 1 | 24 |
| | Head & Neck | Thyroidectomy (any) | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 |
| | Parathyroidectomy (any) | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Branchial cleft cyst/sinus | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Thyroglossal duct cyst/sinus | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Cystic hygroma/lymphangioma | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Dermoid/other cyst | 0.3 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 0 | 0.3 | 1 | 0 | 0 | 3 |
| | Major tumor (head & neck) | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Other (head & neck) | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | TOTAL - Head & Neck | 0.4 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 0 | 0.4 | 1 | 0 | 0 | 3 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Teaching Assistant

[PAGE 2] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|----------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Thoracic | Tracheostomy | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Laryngeal or tracheal resect and/or recon | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Repair chest wall deformity | 0.2 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 3 |
| | Resection chest wall tumor | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Excision mediastinal cyst | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Excision mediastinal tumor | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Pulmonary resection: open | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Pulmonary resection: scope | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Thoracotomy for trauma | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Lung biopsy: open | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Lung biopsy: scope | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Decortication/pleurectomy/blebectomy: open | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Decortication/pleurectomy/blebectomy: scope | 0.2 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 | 0.2 | 0 | 0 | 0 | 1 |
| | Esophageal resection or replacement | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Esophagomyotomy | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Repair esophageal atresia/tracheoesophageal fistula | 0.2 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 1 | 0 | 0 | 3 |
| | Thoracoscopy: other | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 |
| | Thoracotomy: other | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Other (thoracic) | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | TOTAL - Thoracic | 1.1 | 1 | 0 | 1 | 5 | 0.1 | 0 | 0 | 0 | 1 | 1.2 | 2 | 0 | 1 | 5 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Teaching Assistant

[PAGE 3]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|----------------|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Diaphragm | Repair diaphragmatic hernia | 0.2 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 2 |
| | Plication of diaphragm | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Transthoracic and/or retroperitoneal exp for scoliosis | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Other (diaphragm) | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | TOTAL - Diaphragm | 0.3 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.3 | 0 | 0 | 0 | 2 |
| Cardiovascular | Patent ductus arteriosis | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Vascular ring | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Any open heart procedure | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Any closed heart procedure | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Aortopexy | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Renal artery reconstruction | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Peripheral artery reconstruction | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Construction or take down AV fistula/shunt | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Surgical placement/removal central access line | 3.4 | 8 | 0 | 1 | 44 | 0.4 | 1 | 0 | 0 | 3 | 3.8 | 8 | 0 | 1 | 46 |
| | Dialysis access insertion/removal | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Cannulate/ECMO | 0.2 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 2 |
| | Major vessel reconstruction | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Vascular trauma | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Vascular: post-operative exploration | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | TOTAL - Cardiovascular | 3.7 | 8 | 0 | 1 | 45 | 0.4 | 1 | 0 | 0 | 3 | 4.1 | 8 | 0 | 2 | 47 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Teaching Assistant

[PAGE 4] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|-----------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Abdominal | Antireflux procedure: open | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Antireflux procedure: scope | 0.3 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.3 | 1 | 0 | 0 | 2 |
| | Pyloroplasty/gastric resection w/wout vagotomy | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Pyloroplasty/gastric resect w/wout vagotomy - lap | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Any gastrostomy/jejunostomy: open | 0.3 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 4 |
| | Any gastrostomy/jejunostomy: scope | 0.7 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 2 | 0.9 | 1 | 0 | 0 | 4 |
| | Pyloromyotomy: open | 0.4 | 1 | 0 | 0 | 5 | 0.0 | 0 | 0 | 0 | 0 | 0.4 | 1 | 0 | 0 | 5 |
| | Pyloromyotomy: scope | 0.8 | 1 | 0 | 0 | 5 | 0.0 | 0 | 0 | 0 | 0 | 0.8 | 1 | 0 | 0 | 5 |
| | Operation for malrotation | 0.5 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 0 | 0.5 | 1 | 0 | 0 | 3 |
| | Duodenal atresia | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Repair intestinal atresia, stenosis or web | 0.3 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 2 |
| | Intestinal resection: open | 0.7 | 1 | 0 | 0 | 5 | 0.1 | 0 | 0 | 0 | 2 | 0.8 | 1 | 0 | 0 | 7 |
| | Intestinal resection: scope | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Intestinal resection/repair or ostomy for IBD | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Intestinal resect/repair or ostomy for IBD - lap | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 1 |
| | Intestinal resection/repair or ostomy for trauma | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 | 0.2 | 0 | 0 | 0 | 2 |
| | Intestinal resect/repair or ostomy for trauma - lap | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Bowel resection: open | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Bowel resection: scope | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Lap/resec for intussusception-malrotation-volvulus | 0.2 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 | 0.2 | 0 | 0 | 0 | 1 |
| | Ostomy for anorectal malformation: open | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Ostomy for anorectal malformation: laparoscopic | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Teaching Assistant

[PAGE 5] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|-----------|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Abdominal | Ostomy for Hirschsprung's: open | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Ostomy for Hirschsprung's: laparoscopic | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Ostomy for other: open | 0.2 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 2 |
| | Ostomy for other: laparoscopic | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Closure/revision any ostomy/fistula: open | 0.4 | 1 | 0 | 0 | 4 | 0.1 | 0 | 0 | 0 | 1 | 0.4 | 1 | 0 | 0 | 4 |
| | Closure/revision any ostomy/fistula: scope | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Appendectomy: open | 0.5 | 1 | 0 | 0 | 8 | 0.0 | 0 | 0 | 0 | 1 | 0.5 | 1 | 0 | 0 | 8 |
| | Appendectomy: scope | 15.0 | 19 | 0 | 8 | 79 | 0.1 | 0 | 0 | 0 | 1 | 15.1 | 19 | 0 | 9 | 79 |
| | Perineal procedure for imperforate anus | 0.2 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.2 | 1 | 0 | 0 | 3 |
| | Pull through for Hirschsprung's: open | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Pull through for Hirschsprung's: scope | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Pull through for IBD or polyposis: open | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Pull through for IBD or polyposis: scope | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Exploratory lap with or without biopsy: open | 0.9 | 1 | 0 | 1 | 4 | 0.5 | 1 | 0 | 0 | 3 | 1.4 | 2 | 0 | 1 | 6 |
| | Exploratory lap with or without biopsy: lap | 0.2 | 0 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 2 | 0.2 | 1 | 0 | 0 | 2 |
| | Excision of omental/mesenteric cyst | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 2 |
| | Omphalocele (any surgical repair) | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Gastroschisis (any surgical repair) | 0.4 | 1 | 0 | 0 | 3 | 0.0 | 0 | 0 | 0 | 0 | 0.4 | 1 | 0 | 0 | 3 |
| | Resection urachal remnant | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Resection omphalomesenteric duct/cyst | 0.2 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 2 |
| | Excision neuroblastoma/adrenal/other retroperitoneal tumor | 0.2 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.2 | 0 | 0 | 0 | 1 |
| | Excision sacrococcygeal teratoma | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Teaching Assistant

[PAGE 6] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|---------------|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Abdominal | Diagnostic laparoscopy | 0.3 | 1 | 0 | 0 | 3 | 0.6 | 1 | 0 | 0 | 7 | 0.9 | 2 | 0 | 0 | 7 |
| | Adrenal: open | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Adrenal: scope | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Other (abdominal) | 0.7 | 1 | 0 | 0 | 5 | 0.1 | 0 | 0 | 0 | 1 | 0.7 | 1 | 0 | 0 | 6 |
| | TOTAL - Abdominal | 24.3 | 23 | 0 | 19 | 100 | 2.1 | 3 | 0 | 1 | 18 | 26.4 | 24 | 0 | 21 | 105 |
| Hernia Repair | Pediatric repair inguinal hernia > 6 months | 0.9 | 2 | 0 | 0 | 12 | 0.1 | 0 | 0 | 0 | 2 | 0.9 | 2 | 0 | 0 | 12 |
| | Infant repair inguinal hernia < 6 months | 0.5 | 1 | 0 | 0 | 4 | 0.2 | 1 | 0 | 0 | 4 | 0.7 | 1 | 0 | 0 | 8 |
| | Repair umbilical hernia | 0.9 | 2 | 0 | 0 | 9 | 0.1 | 0 | 0 | 0 | 1 | 1.0 | 2 | 0 | 0 | 10 |
| | Repair ventral hernia | 0.2 | 1 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 3 |
| | Repair femoral hernia | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Hernia repair: scope | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 2 |
| | Other (hernia repair) | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | TOTAL - Hernia Repair | 2.6 | 4 | 0 | 1 | 21 | 0.4 | 1 | 0 | 0 | 8 | 3.0 | 5 | 0 | 2 | 24 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Teaching Assistant

[PAGE 7]

Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|---------------|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Liver/Biliary | Major hepatic resection/repair: tumor | 0.1 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 1 |
| | Major hepatic resection/repair: trauma | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Major hepatic resection/repair: other | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Lysis of adhesions | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 2 | 0.1 | 1 | 0 | 0 | 3 |
| | Liver: billiary atresia | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Liver biopsy: open | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 2 |
| | Liver harvest | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Liver transplant | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Cholecystectomy w/wo common bile duct exploration: open | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Cholecystectomy w/wo common bile duct exploration: scope | 1.8 | 3 | 0 | 1 | 18 | 0.1 | 0 | 0 | 0 | 1 | 1.9 | 3 | 0 | 1 | 18 |
| | Portoenterostomy | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Excision choledochal cyst | 0.1 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 1 | 0.2 | 0 | 0 | 0 | 2 |
| | Portosystemic shunts or other operations for portal hypertensior | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Pancreatic resection for: trauma | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Pancreatic resection for: hyperinsulinism | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Pancreatic resection for: tumor | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Pancreas: other | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Operations for pseudocyst | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Splenorrhaphy | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Splenectomy: open | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Splenectomy: scope | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Other (liver) | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | TOTAL - Liver/Biliary | 2.2 | 3 | 0 | 1 | 18 | 0.3 | 1 | 0 | 0 | 3 | 2.5 | 3 | 0 | 1 | 18 |

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Teaching Assistant

[PAGE 8] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|--|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Genito Urinary | Nephrectomy (total or partial): tumor | 0.3 | 1 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.3 | 1 | 0 | 0 | 2 |
| | Nephrectomy (total or partial): trauma | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Nephrectomy (total or partial/nephorrhaphy) | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Nephrectomy (total or partial): other | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Renal transplant | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | Renal: scope | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Renal biopsy (open) | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Cystectomy (partial or total) | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Operation for nephro-uretero lithiasis | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Enteric conduit (any) | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Bladder augmentation or repair (any) | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Pyeloplasty/UPJ reconstruction | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Ureteral reconstruction/reimplantation | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Reconstruct cloacal extrophy | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Circumcision (OR only) | 0.5 | 1 | 0 | 0 | 7 | 0.0 | 0 | 0 | 0 | 1 | 0.5 | 1 | 0 | 0 | 7 |
| | Orchidopexy: open | 0.1 | 0 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 2 | 0.2 | 1 | 0 | 0 | 4 |
| | Orchidopexy: scope | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Orchiectomy | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Operation for torsion testis/appendages | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Operation for varicocele: open | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| Operation for varicocele: scope | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | |
| Procedures for intersex | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | |
| Vaginal procedures: open | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 | |
| Oophorectomy (partial or total) | 0.6 | 2 | 0 | 0 | 8 | 0.0 | 0 | 0 | 0 | 0 | 0.6 | 2 | 0 | 0 | 8 | |
| Hysterectomy/salpingectomy: open | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | |
| Hysterectomy/salpingectomy: scope | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 | |
| Rpr complex laceration vagina/perineum | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 | |
| Other (genito urinary) | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | |
| TOTAL - Genito Urinary | | 2.0 | 2 | 0 | 1 | 10 | 0.1 | 0 | 0 | 0 | 2 | 2.1 | 3 | 0 | 1 | 10 |

Based upon the number of procedures logged by residents in the indicated role.

PEDIATRIC SURGERY: NATIONAL RESIDENT REPORT (Main Table)
Reporting Period: Total Experience of Residents Completing Programs in 2013-2014
Residency Review Committee for Surgery
Report Date: October 30, 2014
Resident Role = Teaching Assistant

[PAGE 9] Number of Programs in the Nation: 34 Number of Residents in the Nation: 39

| | | Primary | | | | | Secondary | | | | | Total | | | | |
|-----------------------|---|--------------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX | Natl Res AVE | Natl Res STD | Natl Res MIN | Natl Res MED | Natl Res MAX |
| RRC Area | RRC Procedure | | | | | | | | | | | | | | | |
| Endoscopic Procedures | Diagnostic thoracoscopy | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Esophageal dilation | 0.1 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 1 | 0.2 | 0 | 0 | 0 | 2 |
| | Esophagoscopy | 0.2 | 1 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 1 | 0.3 | 1 | 0 | 0 | 3 |
| | Esophagoscopy: gastroscopy | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 |
| | Cystoscopy/colposcopy | 0.0 | 0 | 0 | 0 | 1 | 0.0 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 2 |
| | Laryngoscopy/bronchoscopy | 0.1 | 0 | 0 | 0 | 1 | 0.1 | 0 | 0 | 0 | 2 | 0.1 | 0 | 0 | 0 | 2 |
| | Removal foreign body esophagus or trachea | 0.4 | 1 | 0 | 0 | 5 | 0.0 | 0 | 0 | 0 | 0 | 0.4 | 1 | 0 | 0 | 5 |
| | Colonoscopy | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| | Sigmoidoscopy | 0.1 | 0 | 0 | 0 | 2 | 0.0 | 0 | 0 | 0 | 0 | 0.1 | 0 | 0 | 0 | 2 |
| | | TOTAL - Endoscopic Procedures | 1.0 | 2 | 0 | 0 | 6 | 0.2 | 1 | 0 | 0 | 4 | 1.2 | 2 | 0 | 0 |
| Trauma | Non-operative Trauma | 1.5 | 7 | 0 | 0 | 46 | 0.0 | 0 | 0 | 0 | 0 | 1.5 | 7 | 0 | 0 | 46 |
| Total Procedures | TOTAL - Tracked Codes | 40.7 | 39 | 0 | 33 | 218 | 3.8 | 6 | 0 | 1 | 30 | 44.5 | 42 | 0 | 35 | 234 |
| Non-Tracked | TOTAL - Non-Tracked Codes | 2.6 | 6 | 0 | 1 | 35 | 0.6 | 1 | 0 | 0 | 5 | 3.2 | 7 | 0 | 1 | 40 |

Based upon the number of procedures logged by residents in the indicated role.